
Diffusivity scaling on shear flow

Zhong-Tian Wang^{1,2}, Zhi-Xiong He¹, Jia-Qi Dong¹, Zhan-Hui Wang¹, Shao-Yong Chen², Chang-Jian Tang²

¹Southwestern Institute of Physics, Chengdu, Sichuan, 610041, China

²College of Physics Science and Technology, Sichuan University, Chengdu, Sichuan, 610065, China

Email address:

wangzt@swip.ac.cn (Zhong-Tian Wang)

To cite this article:

Zhong-Tian Wang, Zhi-Xiong He, Jia-Qi Dong, Zhan-Hui Wang, Shao-Yong Chen, Chang-Jian Tang. Diffusivity Scaling on Shear Flow. *American Journal of Modern Physics*. Vol. 3, No. 5, 2014, pp. 202-206. doi: 10.11648/j.ajmp.20140305.12

Abstract: Diffusivity scaling on shear flow is investigated. Radial electrical field is the drive of the flow. The turning points of the trapped particle are not on the drift surface, but modified by the radial electrical field. For the first time, an analytical expression of the banana width in presence of shear flow is accurately derived. The particle diffusivity given by Rosenbluth is reproduced but with the shear flow modification.

Keywords: Tokamak Plasma, Diffusivity Scaling, Shear Flow, Trapped Particle, Guiding-Center

1. Introduction

It is generally claimed that the shear flow plays an important role in the onset of the transition from the L mode to H-mode. Experimental evidence¹ also showed that the plasma rotates rapidly in the improved confinement regime, implying that the radial electric field is generated. Neoclassical ion transport in rotating axisymmetric plasma has been systematically investigated by Hinton². However, the energy loss sometimes in the experiment is smaller than the standard neoclassical prediction. The improvement is attributed to the squeezing factor³ treated as the shear flow in this paper. There are renewed interests in the neoclassical fluxes⁴⁻⁶ inside the thermal barriers. The diffusivity scaling on the shear flow presented by Catto⁶ and Shaing⁵ are different. There are intense arguments⁶ on this topic. It is important because it is related to the role of the shear flow in the thermal barrier. Physicists should give a definite answer. The discrepancy between Catto⁶ and Shaing⁵ comes from the trapped particle dynamics which is emphasized in this paper. Accurate banana width, bounce frequency, and the turning point position in the presence of the shear flow are presented. Drift kinetic equation is solved with a particle-conserved Krook collision operator. Rosenbluth's result⁷ is reproduced but with the shear flow modification. The fraction of the trapped particles estimated here is the same as Shaing's⁵, however, diffusivity scaling on shear flow is different.

In section 2, a set of canonical guiding-center variables is

derived by area-conserved transformation. Dynamics for the trapped particles is given in section 3. The drift kinetic equation is solved and the diffusivity scaling is derived in section 4. Summary is presented in the last section.

2. Canonical Guiding-Center Variables

For a tokamak configuration, the Hamiltonian of a charged particle can be expressed as:

$$H = \frac{1}{2M} [(P_R - eA_R)^2 + (P_Z - eA_Z)^2] + (P_\phi - eRA_\phi)^2 / R^2 + e\Phi, \quad (1)$$

where A_R , A_Z , and A_ϕ are the vector potential components of the magnetic field, Φ the electrical potential assumed to be a function of the poloidal magnetic flux Ψ , M the mass of the charged particle set equal to unity for simplicity, and e the charge. P_R , P_ϕ and P_Z are the canonical momentum in the cylindrical coordinates R , ϕ , and Z , respectively, which are as following:

$$P_R = v_R + eA_R, \quad (2)$$

$$P_\phi = Rv_\phi + eRA_\phi, \quad (3)$$

$$P_Z = v_Z + eA_Z. \quad (4)$$

The magnetic field in tokamaks can be expressed as:

$$B = \nabla \phi \times \nabla \Psi + I \nabla \phi, \quad (5)$$

where I is related to the poloidal current. The vector potential components are

$$A_R = 0, \quad A_Z = -I \ln \frac{R}{R_0}, \quad A_\phi = -\frac{\Psi}{R}. \quad (6)$$

As stated by Littlejohn, there has been a gradual evolution over the years away from the averaging approach and towards the transformation approach⁸. We introduce a generating function⁹ to change the cylindrical variables to guiding-center variables:

$$F_1 = -\frac{\Omega_0 R_0^2}{2} \exp\left(\frac{X}{\Omega_0 R_0}\right) \left(\ln \frac{R}{R_0} - \frac{X}{\Omega_0 R_0}\right)^2 \tan \alpha - ZX \quad (7)$$

where

$$X = \Omega_0 R_0 \ln \frac{R_C}{R_0}, \quad (8)$$

Ω_0 is the toroidal gyrofrequency at the magnetic axis, ρ the Larmor radius, α the gyrophase, and the subscripts 0 and c refer to the values at the magnetic axis and the guiding center, respectively. X and α are the new coordinates conjugate to the new momenta:

$$P_X = Z + \rho \sin \alpha + \frac{\rho^2}{4R_C} \sin 2\alpha, \quad (9)$$

$$P_\alpha = \frac{1}{2} \Omega_C \rho^2, \quad (10)$$

where P_X is the guiding center of the Z coordinate, Z_c . The momentum is often transformed into the coordinate during area-conserved canonical transformation⁹.

The Hamiltonian is rewritten as:

$$H = \Omega_C P_\alpha \left[\left(\frac{R_C}{R}\right)^2 \sin^2 \alpha + \cos^2 \alpha \right] + \frac{1}{2R^2} [P_\phi + e\Psi]^2 + e\Phi. \quad (11)$$

The canonical transformation makes the Hamiltonian exact in the new coordinates. The canonical guiding-center variables, $P_\alpha, P_\phi, P_X, \alpha, \phi, X$, are derived and satisfy the Hamiltonian equations:

$$\dot{P}_i = -\frac{\partial H}{\partial q_i}, \quad (12)$$

$$\dot{q}_i = \frac{\partial H}{\partial P_i}, \quad (13)$$

where the P and q are known as the generalized momenta and coordinates. The Jacobian is unity for the area-conserved transformation⁹, that is,

$$J = dP_\alpha dP_X dP_\phi d\alpha dX d\phi = 1. \quad (14)$$

For the tokamaks, the ordering is

$$\delta \sim \rho / r \sim r / R \sim B_\theta / B_\phi, \quad (15)$$

where r and R are the minor and major radii respectively and $R = R_0 + r \cos \theta$. To the first order, the gyro-averaged Hamiltonian in Eq. (11) is approximately expressed as:

$$H = P_\alpha \Omega_c + \frac{1}{2R^2} (P_\xi + e\Psi_c)^2 + e\Phi_c. \quad (16)$$

We have a set of equations of motion in the guiding-center system:

$$v_\Psi = \frac{B_R}{B_p} v_d, \quad (17)$$

$$v_p = \frac{B_p}{B_\phi} \left(v_\phi - \frac{R v_E}{R_0} \right) + \frac{B_z}{B_p} v_d, \quad (18)$$

where $v_\phi = (P_\phi + e\Psi_c) / R$, $v_E = -R_0 \frac{\partial \Phi}{\partial \Psi}$, and

$v_d = \frac{1}{\Omega_0 R_0} (\Omega_C \mu + v_\phi^2)$, v_Ψ is in the radial direction whereas v_p is

in the poloidal direction. Equations (17) and (18) are the generalized versions of the equations of motion obtained by Balescu¹⁰.

3. Dynamics for the Trapped Particles

For the trapped particles, the toroidal velocity is smaller than the perpendicular velocity,

$$v_\phi / v_\perp = v_\phi / (2\Omega_C P_\alpha)^{\frac{1}{2}} \sim \delta^{\frac{1}{2}}. \quad (19)$$

In the rotation frame, we can construct a Hamiltonian in the developed canonical variables, $P_\alpha, P_\phi, P_X, \alpha, \phi, X$,

$$H = \Omega_0 P_\alpha e^{-\frac{X}{\Omega_0 R_0}} + \frac{1}{2R_0^2 S} e^{-\frac{2X}{\Omega_0 R_0}} \left[P_\phi + e\Psi - R_0 v_E e^{\frac{2X}{\Omega_0 R_0}} \right]^2, \quad (20)$$

$$= \frac{1}{2} u_\perp^2 + \frac{1}{2} u_\phi^2$$

where $u_\perp = v_\perp$, $u_\phi = \frac{P_\phi + e\Psi - R v_E}{S^{\frac{1}{2}} R} \approx \frac{v_\phi}{S^{\frac{1}{2}}}$, $S = 1 + \frac{\rho_p^2 e\Phi''(r)}{2T_i}$,

ρ_p is the poloidal Larmor radius, v_E / v_t is the order of δ , $v_t = \sqrt{2T / M}$ is the particle thermal velocity, S is the squeezing factor, the shear flow. The different forms of the Hamiltonian describe the same motion. Eqs.(17) and (18) can be reproduced from Eq.(20).

With the small inverse aspect ratio approximation, we have

$$u_\phi = u_{\phi 0} \sqrt{1 - k^2 \sin^2 \frac{\theta}{2}}, \quad (21)$$

$$u_{\phi 0} = [2H - 2\Omega_0 P_\alpha (1 - \frac{r_0}{R_0})]^{\frac{1}{2}}, \quad (22)$$

$$k = (\frac{2\mathcal{E}u_\perp^2}{u_{\phi 0}^2})^{\frac{1}{2}}. \quad (23)$$

The r_0 should be banana center position. The turning points θ_i of the banana orbit are decided by the following formula,

$$k^2 \sin^2 \frac{\theta}{2} = 1. \quad (24)$$

Once $\theta_i = \pi$ we have $k^2 = 1$ and $u_{\phi \max} = (2\mathcal{E}u_\perp^2)^{\frac{1}{2}}$ for the trapped particles.

We set Ψ_0 as the banana center surface where turning points are evaluated and then we expand Ψ and v_E near Ψ_0 . The banana width and the position of banana center surface are obtained after the expansion,

$$u_\phi = \frac{P_\phi + e\Psi - Rv_E}{S^{\frac{1}{2}}R} = \frac{P_\phi + e\Psi_0 - Rv_{E0} + SR\Omega_p\Delta}{S^{\frac{1}{2}}R}, \quad (25)$$

where Δ is the banana width,

$$\Delta = \frac{u_\phi}{S^{\frac{1}{2}}\Omega_p} = \frac{u_{\phi 0}}{S^{\frac{1}{2}}\Omega_p} \sqrt{1 - k^2 \sin^2 \frac{\theta}{2}}, \quad (26)$$

$$\Psi_0 = \Psi_* + Rv_{E0}/e, \quad (27)$$

where $\Psi_* = -P_\phi/e$ is drift surface and v_{E0} is the value of v_E on the banana center surface. We can see that the turning points of the trapped particle are not on the drift surface but shifted to banana center surface due to the radial electrical field.

We form an invariant⁹ variable,

$$\Pi = \frac{1}{2\pi} \oint P_X dX = \frac{\Omega_0}{2\pi} \oint Z_c \frac{R_0}{R_c} dR_c \approx \frac{\Omega_0}{2\pi} \oint r_0 \Delta d\theta, \quad (28)$$

which is actually the flux enclosed by banana orbit. Now we introduce a new angle which satisfies the equation,

$$\sin^2 \beta = k^2 \sin^2 \frac{\theta}{2}, \quad (29)$$

then,

$$\begin{aligned} \Pi &\approx \frac{\Omega_0 r_0 u_{\phi 0}}{\pi S^{\frac{1}{2}} \Omega_p} \int_{-\theta_i}^{\theta_i} \sqrt{1 - k^2 \sin^2 \frac{\theta}{2}} d\theta = \frac{4qR_0 u_{\phi 0} k_1}{\pi S^{\frac{1}{2}}} \int_0^{\frac{\pi}{2}} \frac{(1 - \sin^2 \beta) d\beta}{\sqrt{1 - k_1^2 \sin^2 \beta}}, \quad (30) \\ &= \frac{8qR_0 (e\Omega_0 P_\alpha)^{\frac{1}{2}}}{\pi S^{\frac{1}{2}}} [E(k_1) - (1 - k_1^2)K(k_1)] \end{aligned}$$

where $k_1^2 = k^{-2}$, K and E are complete elliptic functions. The bounce frequency of the trapped particle is

$$\omega_b = \frac{\partial H}{\partial \Pi} = \frac{(2\mathcal{E}Sv_\perp^2)^{\frac{1}{2}}}{2qR_0} \left(\frac{\pi}{2K(k_1)} \right). \quad (31)$$

Eqs. (21-31) give, for the first time, a so clear picture of the trapped particle motion dependent on the shear flow. It is real motion no other choice.

For a Maxwellian distribution function, the fraction of the trapped particles can easily calculated via

$$F = 2\sqrt{\pi} \int_0^\infty y dy \int_{-x_{\max}}^{x_{\max}} e^{-y^2 - x^2} dx, \quad (32)$$

where $y = v_\perp / v_t$, $x = v_\parallel / v_t$, and

$v_{\phi \max} = S^{\frac{1}{2}} u_{\phi \max} = S^{\frac{1}{2}} (2\mathcal{E}v_\perp^2)^{\frac{1}{2}}$. For the trapped particles, x is small, thus it leads

$$F = 4\sqrt{\pi} \int_0^\infty dy e^{-y^2} y^2 (2\mathcal{E}S)^{\frac{1}{2}} = (2\mathcal{E}S)^{\frac{1}{2}}, \quad (33)$$

which agrees with Shaing's⁵ and disagrees with Catto's⁶. Since the trapped particle pitch angle is limited, the effective collision frequency ν_{eff} here should be $\nu / (u_\phi^2 / v^2)$. Thus,

the diffusivity is $F\Delta^2 \nu v^2 / u_\phi^2 \approx \nu \rho_p^2 (2\mathcal{E}/S)^{\frac{1}{2}}$ which is different with Shaing's⁵.

4. Neoclassical Transport

To illustrate the significance of the trapped-particle dynamics, the particle diffusivity is calculated. Instead of canonical gyrokinetic variables, $P_\alpha, P_\phi, P_X, \alpha, \phi, X$, we use extended phase space variables¹¹, $P_\alpha, P_\phi, r, H, \alpha, \phi, \beta, t$, in the drift kinetic equation,

$$\frac{\partial f}{\partial t} + \frac{d\beta}{dt} \frac{\partial f}{\partial \beta} + \frac{dr}{dt} \frac{\partial f}{\partial r} = C(f). \quad (34)$$

We define an averaged angle velocity,

$$\left\langle \frac{d\beta}{dt} \right\rangle = \frac{1}{T} \oint \frac{d\beta}{dt} dt = \frac{2\pi}{T}, \quad (35)$$

where T is the bounce period. We can see that the averaged angle velocity is the bounce frequency in Eq. (31). To illustrate flow shear effects on the neoclassical transport, we take

$$f = f_* + g = f_* + g_0 + g_{1s} \sin \beta + g_{1c} \cos \beta, \quad (36)$$

where f_* is equilibrium distribution function in a Maxwellian form with H at the place of energy and P_ϕ at the place of position,

$$f_* = F_m(r) \left[1 - \frac{v_\phi}{\Omega p} \frac{\partial \ln n}{\partial r} \right], \quad (37)$$

$$\frac{\partial f_*}{\partial r} = 0. \quad (38)$$

To catch the key points and avoid the complexity we only consider density gradient. Temperature gradient can change the expression of diffusivity, but can not change the scaling on the shear flow.

Eq. (34) is rewritten as

$$\frac{\partial \beta}{\partial t} \frac{\partial g}{\partial \beta} - \sin \theta v_d \frac{\partial g}{\partial r} = C(g - g_m) - \frac{\partial f}{\partial t}, \quad (39)$$

$$g_m = \frac{v_\phi}{\Omega_p} F_m \frac{\partial \ln n}{\partial r}. \quad (40)$$

To make Eq. (39) tractable, we take

$$\frac{d\beta}{dt} \approx \frac{d\beta}{dt} = \omega_b, \quad (41)$$

and

$$C(g - g_m) = \nu_{eff} (g - g_m). \quad (42)$$

Furthermore, it is assumed that k_1 and θ_i are small, which means deeply trapped particles dominate. In Eq. (36) g_{1s} and g_{1c} are one order smaller than g_0 . From Eq. (29) and

$$\text{Eq. (39) we obtain } g_0 = g_m, \quad g_{1s} = \frac{2k_1 \nu_{eff} \frac{\partial g_0}{\partial r}}{\omega_b^2 + \nu_{eff}^2}, \quad g_{1c} = -\frac{\omega_b}{\nu_{eff}}.$$

With Eq. (37) and Eq. (38) the drift equation of Eq. (39) after the bounce average turns to be

$$\frac{\partial f}{\partial t} - \frac{\partial}{\partial r} \frac{2\nu_{eff} k_1^2 v_d^2}{\omega_b^2 + \nu_{eff}^2} \frac{\partial F_m}{\partial r} = 0. \quad (43)$$

In the banana regime, we have

$$\frac{\nu_{eff}^2}{\omega_b^2} \approx \delta. \quad (44)$$

After integrating Eq. (43) over velocity space, we get the continuity equation

$$\frac{\partial n}{\partial t} + \frac{\partial}{\partial r} \Gamma = 0, \quad (45)$$

$$\text{where } \Gamma = -D \frac{\partial n}{\partial r},$$

$$D = 2 \int_0^\infty y dy \int_{-x_{\max}}^{x_{\max}} dx \frac{2\nu_{eff} k_1^2 v_d^2}{\sqrt{\pi} \omega_b^2} e^{-y^2 - x^2} = 0.75 \nu \sqrt{2\varepsilon} \rho_p^2 / S^{\frac{1}{2}}, \quad (46)$$

where $x = \frac{v_\phi}{v_t}$, $y = \frac{v_\perp}{v_t}$, $x_{\max} = \frac{\sqrt{2\varepsilon S} v_\perp}{v_t} = \sqrt{2\varepsilon S} y$. From Eq. (46) we can see that Rosenbluth's result⁹ is reproduced but with the shear flow modification.

5. Summary

The area-conserved transformation proposed by Lichtenberg and Lieberman⁹ is employed. A complete set of canonical guiding-center variables, $P_\alpha, P_\phi, P_X, \alpha, \phi, X$, are derived. The accurate relation between the particle motion and the shear flow for the trapped particles is also derived,

$$\text{including the banana width } \Delta = \frac{u_\phi}{S^{\frac{1}{2}} \Omega_p} = \frac{u_{\phi 0}}{S^{\frac{1}{2}} \Omega_p} \sqrt{1 - k^2 \sin^2 \frac{\theta}{2}},$$

the position of banana center surface $\Psi_0 = \Psi_* + R v_{E0} / e$

$$\text{and the bounce frequency } \omega_b = \frac{\partial H}{\partial \Pi} = \frac{(2\varepsilon S v_\perp^2)^{\frac{1}{2}}}{2qR_0} \left(\frac{\pi}{2K(k_1)} \right).$$

For a Maxwellian distribution function, the fraction of the trapped particles is calculated as

$$F = 4\sqrt{\pi} \int_0^\infty dy e^{-y^2} y^2 (2\varepsilon S)^{\frac{1}{2}} = (2\varepsilon S)^{\frac{1}{2}} \quad \text{which agrees with}$$

Shaing's⁵ and disagrees with Catto's⁶. Since the trapped particle pitch angle is limited, the effective collision frequency ν_{eff} here should be $\nu / (u_\phi^2 / v^2)$, therefore, the

diffusivity is $F \Delta^2 \nu v^2 / u_\phi^2 \approx \nu \rho_p^2 (2\varepsilon / S)^{\frac{1}{2}}$ which is different with Shaing's⁵. Drift kinetic equation is solved with particle-conserved Krook collision operator. Rosenbluth's result⁷ is reproduced but with shear flow modification in this paper.

Acknowledgments

It is appreciated that Dr. R. D. Hazeltine and Dr. P. Morrison checked and verified the generating function when author Z. T. Wang worked at IFS of the University of Texas.

This work was supported by Natural Science Fund No. 11261140327, No.11205053, No.11005035, No.11105046 and The National Magnetic Confinement Fusion Science Program (Grant No. 2013GB107002).

References

- [1] E.J.Doyle, R.J. Groebner, K.M.Burrell, P. Gohil, T. Lehcka, N. C. Luhmann Jr., H. Matsumoto, T. H. Osborne, W. A. Peebles, and R. Philipona, Modification in turbulence and edge electric field at the L-H transition in DIII-D tokamak, *Phys. Fluids B* 3, 2300(1991).
- [2] F.L.Hinton and S.K.Wong, Neoclassical ion transport in rotating axisymmetrical plasmas, *Phys. Fluids* 28, 3082(1985).
- [3] R.D.Hazeltine, Self-consistent radial sheath, *Phys.Fluids B*1, 2031(1989).
- [4] G. Kagan and P. J. Catto, *Phys. Rev. Lett.* 105, 045002(2010).
- [5] K. C. Shaing and C. T. Hsu, *Phys. Plasmas* 19, 022502(2012).
- [6] P. J. Catto, F. I. Parra, G. Kagan, J. B. Parker, I. Pusztai, and M. Landreman, *Plasma Phys. Control. Fusion*, 55, 045009(2013).
- [7] M. N. Rosenbluth, R. D. Hazeltine, F. L. Hinton, *Phys. Fluids*, 15, 116(1972).
- [8] R. G. Littlejohn, *J. Plasma physics* 29, 111(1983).
- [9] A. J. Lichtenberg and Lieberman, *Regular and Stochastic Motion*, *Applied Sciences* 38, (Springer-Verlag New York Inc. 1983).
- [10] R. Balescu, in *Transport Processes in Plasma*, (North-Holland, Amsterdam · Oxford · New York · Tokyo, 1988), Vol.2, P.393.