

Analysis of the Determinants of Female Genital Mutilation Practices in Senegal: A Secondary Analysis of the 2018 DHS

Ndeye Mareme Sougou^{1,2,*}, Jean Baptiste Niokhor Diouf³, Oumar Bassoum^{1,2}, Ibrahima Seck¹

¹Department of Preventive Medicine and Public Health, Faculty Medicine, University Cheikh Anta Diop, Dakar, Senegal

²Institute of Health and Development, University Cheikh Anta Diop, Dakar, Senegal

³Hospital Roi Baudouin, Faculty Medicine, University Iba Mar Diop, Dakar, Senegal

Email address:

nmsougou@hotmail.com (N. M. Sougou), jeanniokhor@gmail.com (J. B. N. Diouf), bassoum.oumar@gmail.com (O. Bassoum),

ibouseck@yahoo.fr (I. Seck)

*Corresponding author

To cite this article:

Ndeye Mareme Sougou, Jean Baptiste Niokhor Diouf, Oumar Bassoum, Ibrahima Seck. Analysis of the Determinants of Female Genital Mutilation Practices in Senegal: A Secondary Analysis of the 2018 DHS. *Central African Journal of Public Health*.

Vol. 7, No. 3, 2021, pp. 121-126. doi: 10.11648/j.cajph.20210703.15

Received: April 30, 2021; Accepted: May 21, 2021; Published: May 31, 2021

Abstract: Introduction: Female genital mutilation (FGM), also known as female genital cutting or circumcision, threatens the health and well-being of millions of girls, women, and their children around the world. In Senegal, FGM practices persist despite numerous interventions. This study aims to analyze the determinants of FGM in Senegal. Methods: This article uses data from the 2018 Senegal Demographic and Health Survey (DHS), a nationally representative survey of 9413 women aged 15-49 years. In the descriptive analysis, variables were presented in terms of frequency and percentage of data. The significance level was set at 5, and 95% confidence intervals (CIs) were used. Variables with p values less than 0.25 in the bivariate analysis were selected for multivariate analysis. The dependent variable was a composite variable generated from 3 DHS variables that described the types of post-FGM genital lesions that exist in women. The analysis was performed using STATA/SE 17. Results: The prevalence of FGM is 17.14%. Eleven-point fifty-two percent (11.52%) or 999 women believe that FGM is a religiously recommended practice. Ninety-nine percent (993) of these women were Muslim ($p=0.0017$). Fifteen-point thirty-nine percent (15.39%) or 1,334 women think it is a practice that should continue to be practiced and 80.59% (or 6,988 women) think it should be stopped. The protective factors for the occurrence of FGM were female empowerment (high level of education of the woman (primary $ajOR=0.64$ [0.50-0.83] and secondary $ajOR=0.43$ [0.32, 0.57]) and the fact that the head of the family was a woman $ajOR$ (0.75 [0.59-0.97]); belonging to the central region of Senegal (Diourbel, Kaolack, Thiès, Louga and Fatick) and Christian religion ($ajOR=0.05$ [0.02-0.13]). The risk factors for female genital mutilation in Senegal were ethnicity and belonging to certain regions in the northeast and southeast of Senegal (Tambacounda, Matam, Kedougou). Conclusion: This study showed that FGM practices are still persistent in Senegal. This study underlined that the empowerment of women would allow the fight against FGM. Health interventions should be multisectoral, involving the education sectors with a strong investment in girls' education and retention.

Keywords: Female Genital Cutting, Determinants, Senegal

1. Introduction

Female genital mutilation (FGM), also known as female circumcision, threatens the health and well-being of millions of girls, women and their children around the world [1]. Female genital mutilation has no health benefits and is known to cause serious short- and long-term damage to the physical and

psychological health of victims [2]. Thus they are at the origin of obstetrical consequences which can engage the vital prognosis of the woman during the delivery [3]. Over 200 million girls and women alive today have been cut in 30 countries in Africa, the Middle East and Asia where FGM is concentrated [4]. Africa still bears the greatest burden. Female genital mutilation of any kind has been recognized as a harmful practice and a violation of the human rights of girls

and women [5]. By 2030, the Sustainable Development Goals through Goal 5 in its indicator 5.3.2 addresses the problem of FGM by insisting on a reduction of the "Proportion of girls and women aged 15-49 years who have undergone female genital mutilation/cutting, by age" [6]. In Senegal, since the 1970s, excision has been combated by the Senegalese authorities through awareness programs. Since 1999, the practice is legally repressed and excisers are regularly brought to justice and sentenced to a minimum of 6 months in prison, at least for example [7]. However, it is noted that over the years, despite the many health interventions made, there is a slight decline in FGM practices in Senegal [7].

Studies have been conducted on FGM in Senegal, but few have focused on the determinants of FGM practices. This study aims to analyze the determinants of FGM in Senegal.

2. Methods

2.1. Type of Study

This is a descriptive cross-sectional study with an analytical purpose.

2.2. Data

This article uses data from the 2018 Senegal Demographic and Health Survey (DHS), a nationally representative survey of 9413 women aged 15 to 49 years. The DHS-Continuous 2018 sample is representative at the national level, at the regional level, and for urban and rural settings. The sample was drawn stratum by stratum. Thus, the sample is based on a stratified, two-stage area sample drawn in accordance with the DHS sampling methodology [8]. At the first level, 400 clusters (primary sampling units [PSUs]) were drawn from the list of enumeration areas (EAs) established during the 2013 General Census of Population and Housing, Agriculture, and Livestock (RGPHAE), using a systematic draw with probability proportional to size, with the PSU size corresponding to the number of households [9]. A count of households in each of these clusters provided a list of households from which a sample of 22 households per cluster was drawn, in the second stage, from both urban and rural areas with systematic equal probability. A total of 8,800 households (4,092 urban and 4,708 rural) were selected.

Data were collected using four questionnaires - Household, Female, Male and Biomarkers. The Female Individual Questionnaire was used to record information from females aged 15-49 years who were residents or visitors the night before the survey. The Individual Record (IR) file was used for the analysis.

The dependent variable is a composite variable generated from 3 DHS variables that describe the types of post-FGM genital lesions that exist in the woman interviewed. The lesion was described by the woman and is not observed by a health provider. These variables are: variable 1: women who have parts of flesh removed from the genital area coded in 3

modalities yes, no, don't know; variable 2: women who have lesions in the genital area that is just nicked without removing flesh with 3 modalities yes, no, don't know; and variable 3: women who have closure of the genital area sewn with 3 modalities yes, no, don't know. The generated variable is dichotomous with 2 modalities "yes" and "no". "Yes" for all women presenting at least one of the above-mentioned lesions.

The independent variables: the socio-demographic factors were place of residence: this was dichotomized as "urban" or "rural"; the woman's level of education; the gender of the head of the household; the household's level of wealth: the wealth index, a measure of relative economic well-being based on household assets, was categorized into quintiles (lowest, second, middle, fourth, highest) and derived from the wealth score, religion, ethnicity, and region of residence.

2.3. Statistical Analysis

The analysis was performed using STATA/SE 17 software. As mentioned above in the data source section, a two-stage sampling design was adopted. To account for the multi-stage sampling design of the survey, all data were weighted to account for disproportionate sampling and non-response. In the descriptive analysis, variables were presented in terms of frequency and percentage of data. Comparisons between groups were made using the Chi 2 test. The significance level was set at 5, and 95% confidence intervals (CIs) were used. Variables with p values less than 0.25 in the bivariate analysis were selected for multivariate analysis [10]. To assess the factors associated with the existence of female genital mutilation, a multivariate logistic analysis was performed to adjust for the effect of confounders. Adjusted odds ratios (ORa) were calculated with their 95% confidence intervals. To manage complex sampling (multistage sampling, weighting, and stratification), variables for identifying weights, strata, and primary sampling units (PSUs) were defined before using SVY (STATA survey prefix).

3. Ethical Approval

This study is a secondary analysis of data from the DHS conducted in Senegal in 2018. The 2018 Demographic and Health Survey (DHS) in Senegal was approved by the National Ethics Committee (NEC). The survey was also approved by ICF's Institutional Review Board. Informed consent was obtained from all participants in writing.

4. Results

Prevalence of FGM

The prevalence of FGM is 17.14% (see Table 1). Most of the women have a mutilation of a part of the reproductive system with 60%, 29.39% of women have Genital area just nicked without removing any flesh et 6.88% des femmes ont une infibulation (voir table 1).

Table 1. Prévalence des MGF au Sénégal en 2018.

Female Genital Mutilation	Frequency n=9414	Percentage%
Yes	1617	17.18
No	7797	82.82
Type de FGM		
Flesh removed from genital area	n=2195	
yes	13 38	60.96
no	566.	25.80
don't know	291.	13.24
Genital area just nicked without removing any flesh	N=857	
yes	252.	29.39
no	333.	38.79
don't know	273.	31.81
Genital area sewn closed	n=2196.	
yes	151.	6.88
no	1838.	83.70
don't know	207.	9.41

Table 2. Distribution of FGM according to socio-demographic characteristics.

	Frequency n=1617	Percentage%
Woman highest educational level		
no education	856	9.097
primary	360	3.823
secondary	352	3.74
higher	49	0.5158
Wealth index		
poorest	452	4.8
poorer	283	3.004
middle	287	3.052
richer	324	3.438
richest	272	2.882
Region		
Dakar	442	4.7
ziguinchor	99	1.05
diourbel	99	0.0761
saint-louis	7	1.468
tambacounda	138	2.009
kaolack	189	0.9057
thies	85	1.02
louga	11	0.1162
fatick	12	0.1314
kolda	156	1.662
matam	189	2.012
kaffrine	17.	0.1839
kedougou	56	0.594
sedhiou	118	1.256
Type of place residence		
rural	820	8.707
urban	797	8.468
Religion		
muslim	1 605	17.05
chistiane	10	0.1074
no religion	0.8	0.0087
other	0.5	0.0052
Ethnicity		
wolof	19	0.1792
poullar	893	9.487
serer	4	0.0462
mandingue/ soce	250	2.661
diola	134	1.424
soninke	92	0.977
senegalese ethnicity	109	1.157
not senegalese	117	1.244
sex of household head		
male	1 205	12.8
female	412	4.378

Distribution of FGM according to socio-demographic characteristics

The prevalence of FGM is higher among women without education (9%). The regions with the highest prevalence are Dakar (4.7%), Tambacounda (2%), Sedhiou (1.2%) and Matam (2%). Women of the Muslim religion are the most concerned (17.05%). The Poular ethnic group is the most represented (9.49%). FGM is more common among women living in a house where the head of the household is a man (12.8%). (see Table 2).

Women's knowledge and opinions about FGM

Women who had heard of FGM represented 90.66% (8,535) of the women.

Eleven-point fifty-two percent (11.52%) or 999 women thought that FGM was a religiously recommended practice. Ninety-nine percent (993) of these women were Muslim ($p=0.0017$).

Fifteen-point thirty-nine percent (15.39%) or 1,334 of the women thought that it was a practice that should continue to be performed and 80.59% (6,988 women) thought that it should be stopped.

Table 3. Factors associated with female genital mutilation in Senegal.

FGM	Odds ratio	[95% conf. interval]	P>t
Woman highest educational level			
No education	1		
Primary*	0.64	[0.50-0.82]	0.001
Secondary*	0.43	[0.32-0.57]	0.000
higher	0.54	[0.24-1.18]	0.123
Wealth index			
Poorest	1		
poorer	0.92	[0.67-1.25]	0.593
middle	0.90	[0.63-1.28]	0.564
richer	0.87	[0.55-1.36]	0.529
richest	0.68	[0.36-1.32]	0.255
Region			
Dakar	1		
Ziguinchor	1.16	[0.63-2.14]	0.622
Diourbel*	0.05	[0.02-0.14]	0.000
Saint-louis	0.81	[0.43-1.50]	0.494
Tambacounda*	2.31	[1.39-3.82]	0.001
Kaolack*	0.38	[0.18-0.78]	0.009
Thies*	0.45	[0.25-0.83]	0.011
Louga*	0.07	[0.03-0.17]	0.000
Fatick*	0.27	[0.13-0.54]	0.000
kolda	0.88	[0.53-1.44]	0.604
Matam*	1.97	[1.17-3.32]	0.011
Kaffrine*	0.25	[0.11-0.58]	0.001
Kedougou*	2.76	[1.42-5.39]	0.003
Sedhiou	1.16	[0.67-2.01]	0.600
Type of place residence			
Urban	1		
Rural	0.87	[0.62-1.22]	0.429
Religion			
Muslim	1		
Christiane*	0.06	[0.02-0.14]	0.000
no religion	1		
Ethnicity			
Wolof	1		
Poular*	83.22	[34.09-203.11]	0.000
serer	0.76	[0.22-2.64]	0.668
mandingue/ soce*	120.27	[46.12-313.67]	0.000
Diola*	114.37	[44.58-293.39]	0.000
Soninke*	85.05	[30.07-240.53]	0.000
senegalese ethnicity*	65.47	[24.72-173.36]	0.000
not senegalese*	141.09	[49.35-403.39]	0.000
sex of household head			
male	1		
Female*	0.76	[0.59-0.97]	0.029

Factors associated with FGM

Protective factors for the occurrence of FGM were women's empowerment factors (high level of education of the woman (primary $ajOR=0.64$ [0.50-0.83] and secondary $ajOR=0.43$ [0.32, 0.57]) and the fact that the head of the family was a woman $ajOR$ (0.75 [0.59-0.97]); belonging to the central region of Senegal (Diourbel, Kaolack, Thiès, Louga and Fatick) and Christian religion ($ajOR=0.05$ [0.02-0.13]). Risk factors for female genital mutilation in Senegal were ethnicity and belonging to certain regions in the northeast and southeast of Senegal (Tambacounda, Matam, Kedougou). (See Table 3).

5. Discussion

It is estimated that more than 200 million girls and women alive today have undergone female genital mutilation in countries where the practice is concentrated. In addition, an estimated 3 million girls are at risk of undergoing female genital mutilation each year [11]. In Senegal, the prevalence rate of FGM is 17.14% in 2018. This high prevalence shown by our study is similar to other countries in Africa. In West Africa is 25.4% in 2017 [12]. For example, in Sudan the prevalence is 20%, 21.1% in Ghana [13, 14].

According to WHO, FGM/C is defined as "all procedures involving partial or total removal of the external female genitalia or other injury to the female genital organs for non-medical reasons." In the medical literature, four main types of FGM/C are recognized [15]:

Type I: excision of the clitoral hood with or without removal of parts or the entire clitoris (clitoridectomy);

Type II: excision of the clitoris together with parts or all of the labia minora;

Type III: excision of parts or the whole of the clitoris, labia minora and labia majora and stitching or narrowing of the introitus, with a very small outlet for passage of urine and menstruum. This is also known as infibulation;

Type IV: other harmful procedures to the female genitalia for non-medical purposes. Examples are—pricking, piercing, incising, scraping and cauterization. Others are hymenectomy, cutting of the vagina and introduction of corrosive substances or herbs into the vagina to cause bleeding or to tighten or narrow the vagina.

In Senegal, most women present as lesion flesh removed from genital area avec 60.96% which corresponds to type I and type II, 29.39% of women have Genital area just nicked without removing any flesh (Type IV) and 6.88% of women have an infibulation (Type III). The type of lesions identified by the DHS cannot be precise because it is based on a description of their anatomy by the respondents and not on direct observation by a health professional [16]. Depending on the type of mutilation, the genito-obstetrical complications are more or less serious but also the reconstruction is less possible [17]. In other countries of the world the most widespread types are type I mutilations, type III being found very rarely [1, 18].

Women who had heard of FGM represented 90.66% (8,535) of women. Fifteen-point thirty-nine percent (15.39%) or 1,334 of the women think that it is a practice that should continue to

be practiced and 80.59% (6,988 women) think that it should be stopped. Numerous health interventions and campaigns have been carried out in African countries to fight against FGM. These policies have led to a decline in the practice but above all to an increase in community awareness of these harmful practices [19]. This situation is not similar to that of countries in the sub-region where women still say they are not yet ready to abandon these practices [20]. Other studies have shown that information dissemination improves cognitions about FGM, but that the success of interventions depends on a range of other contextual factors [21].

Eleven-point fifty-two percent (11.52%) or 999 women believe that FGM is a religiously recommended practice. Ninety-nine percent (993) of these women were Muslim ($p=0.0017$). This is also the case in other countries where the prevalence of FGM is high and where one of the main reasons for the persistence of this practice is that communities believe that FGM is recommended by religion [19, 22]. In Senegal, in some religious communities of the Pular ethnic group, despite laws that condemn these practices, they continue to be legitimized by religious arguments [23]. Our study showed that religion is indeed a factor associated with FGM practices.

The analysis of factors associated with FGM also showed that region of residence was a determinant of FGM. It was found that some regions, such as the Southeast, are strongly associated with FGM.

In our study, the protective factors for FGM are factors of women's empowerment. Thus, a high level of education would protect women from these practices (primary level $oraj$: 0.64 [0.50-0.82]; secondary level $oraj$: 0.43 [0.32-0.57]). Other studies had shown that FGM was less prevalent among educated women (p value < 0.001) [13, 18]. Empowerment factors were described as necessary to work on reducing FGM practices [24]. In the same register, our study showed that households headed by women were less likely to undergo FGM ($ORaj$: 0.76 [0.59-0.97]). In Guinea, the factors of empowerment of women through the strengthening of decision-making power allowed them to stand up against FGM practices [25]. Empowerment of women and girls was seen as an essential intervention to eradicate this practice [26].

Limitations

This study has some limitations. In particular, the analyses used cross-sectional data, so only associations and not causal relationships were established. This study could be complemented by a qualitative study to understand the contextual and socio-cultural factors that cause FGM practices to persist despite all the interventions implemented.

6. Conclusion

This study has shown that FGM practices are still persistent in Senegal. They are mostly practiced in certain regions of Senegal and by certain ethnic groups. Religious justification is still prevalent in the persistence of FGM. This study has highlighted that the empowerment of women would allow the fight against FGM. Health interventions should be multisectoral by involving the education sectors with a strong

investment in girls' education and their retention in school. Households headed by women suffer less from this practice. This result shows the importance of women's decision-making autonomy, which could be an important lever on which policies could rely in the fight against FGM. In the future, this study could be complemented by qualitative studies that could explore the socio-cultural norms and social processes at work in these specific ethnic groups and regions of Senegal where high prevalence of FGM persists.

References

- [1] Koski A, Heymann J. Thirty-year trends in the prevalence and severity of female genital mutilation: a comparison of 22 countries. *BMJ Global Health*. 8.
- [2] Reisel D, Creighton SM. Long term health consequences of Female Genital Mutilation (FGM). *Maturitas*. janv 2015; 80 (1): 48-51.
- [3] Berg RC, Underland V. The obstetric consequences of female genital mutilation/cutting: a systematic review and meta-analysis. *Obstet Gynecol Int*. 2013; 2013: 496564.
- [4] Female genital mutilation [Internet]. [cité 26 avr 2021]. Disponible sur: <https://www.who.int/news-room/fact-sheets/detail/female-genital-mutilation>.
- [5] Weltgesundheits organisation, éditeur. Eliminating female genital mutilation: an interagency statement; OHCHR, UNAIDS, UNDP, UNECA, UNESCO, UNFPA, UNHCR, UNIFEM, WHO. Geneva: World Health Organization; 2008. 40 p.
- [6] SDG Indicators — SDG Indicators [Internet]. [cité 26 avr 2021]. Disponible sur: <https://unstats.un.org/sdgs/metadata/?Text=&Goal=5&Target=5.3>.
- [7] Les mutilations sexuelles féminines (MSF). Division de l'Information, de la Documentation et des Recherches de l'Ofpra. 2019, 15p [Internet]. [cité 26 avr 2021]. Disponible sur: https://ofpra.gouv.fr/sites/default/files/atoms/files/1905_sen_mgf.pdf
- [8] ICF International. 2012. Demographic and Health Survey Sampling and Household Listing Manual. MEASURE DHS, Calverton, Maryland, U.S.A.: ICF International. 98p.
- [9] Croft, Trevor N., Aileen M. J. Marshall, Courtney K. Allen, et al. 2018. Guide to DHS Statistics. Rockville, Maryland, USA: ICF <https://dhsprogram.com/data/Guide-to-DHS-Statistics/index.cfm>.
- [10] Collett D. Modelling Binary Data, Second Edition. CRC Press; 2002. 406 p.
- [11] Prevalence of female genital mutilation [Internet]. [cité 28 avr 2021]. Disponible sur: <https://www.who.int/teams/sexual-and-reproductive-health-and-research/areas-of-work/female-genital-mutilation/prevalence-of-female-genital-mutilation>.
- [12] Kandala N-B, Ezejiofor MC, Uthman OA, Komba P. Secular trends in the prevalence of female genital mutilation/cutting among girls: a systematic analysis. *BMJ Global Health*. 1 oct 2018; 3 (5): e000549.
- [13] Satti A, Elmusharaf S, Bedri H, Idris T, Hashim MSK, Suliman GI, et al. Prevalence and determinants of the practice of genital mutilation of girls in Khartoum, Sudan. *Annals of Tropical Paediatrics*. 1 déc 2006; 26 (4): 303-10.
- [14] Oduro AR, Ansah P, Hodgson A, Afful TM, Baiden F, Adongo P, et al. Trends in the prevalence of female genital mutilation and its effect on delivery outcomes in the kassena-nankana district of northern Ghana. *Ghana Medical Journal* [Internet]. 2006 [cité 28 avr 2021]; 40 (3). Disponible sur: <https://www.ajol.info/index.php/gmj/article/view/55258>.
- [15] Odukogbe A-TA, Afolabi BB, Bello OO, Adeyanju AS. Female genital mutilation/cutting in Africa. *Transl Androl Urol*. 2017; 6 (2): 138-48.
- [16] Sénégal - 2018_01.pdf [Internet]. [cité 28 avr 2021]. Disponible sur: https://www.rhsupplies.org/uploads/tx_rhscpublications/S%C3%A9n%C3%A9gal_-2018_01.pdf.
- [17] Puppo V. Female genital mutilation and cutting: An anatomical review and alternative rites. *Clin Anat*. janv 2017; 30 (1): 81-8.
- [18] Al Awar S, Al-Jefout M, Osman N, Balayah Z, Al Kindi N, Ucenic T. Prevalence, knowledge, attitude and practices of female genital mutilation and cutting (FGM/C) among United Arab Emirates population. *BMC Womens Health*. 22 avr 2020; 20 (1): 79.
- [19] Awolola OO, Ilupeju NA. Female genital mutilation; culture, religion, and medicalization, where do we direct our searchlights for its eradication: Nigeria as a case study. *Tzu Chi Med J*. 2019; 31 (1): 1-4.
- [20] Dicko-Traoré F, Diakité FL, Diakité AA, Konaté D, Keïta JT, Traoré F, et al. Connaissances, attitudes et pratiques des mères relatives à l'excision à bamako. *Mali Med*. 31 mars 2014; 30-5.
- [21] Berg RC, Denison E. Interventions to reduce the prevalence of female genital mutilation/cutting in African countries. *Campbell Systematic Reviews*. 2012; 8 (1): 1-155.
- [22] Shabila NP, Ahmed HM, Safari K. Knowledge, attitude, and experience of health professionals of female genital mutilation (FGM): A qualitative study in Iraqi Kurdistan Region. *Health Care Women Int*. nov 2017; 38 (11): 1202-18.
- [23] Sylla PS. Interdiction de l'excision au Sénégal: pour comprendre les raisons de l'inefficacité de la loi. Centre Interdisciplinaire d'Études et de Recherche de la Vallée (CIERVAL), Université Gaston Berger. 2002; 18p.
- [24] Diop NJ, Askew I. The Effectiveness of a Community-based Education Program on Abandoning Female Genital Mutilation/Cutting in Senegal. *Studies in Family Planning*. 2009; 40 (4): 307-18.
- [25] Doucet M-H, Delamou A, Manet H, Groleau D. Beyond will: the empowerment conditions needed to abandon female genital mutilation in Conakry (Guinea), a focused ethnography. *Reprod Health*. déc 2020; 17 (1): 61.
- [26] Williams-Breault BD. Eradicating Female Genital Mutilation/Cutting. *Health Hum Rights*. déc 2018; 20 (2): 223-33.