

(Alpha, Beta)-Normal and Skew n-Normal Composite Multiplication Operator on Hilbert Spaces

Senthil^{1,*}, Nithya², Suryadevi³, David Chandrakumar³

¹Department of Economics and Statistics, Government of Tamilnadu, DRDA, Dindigul, India

²Department of Mathematics, Mother Teresa Women's University, Kodaikanal, India

³Department of Mathematics, Vickram College of Engineering, Enathi, India

Email address:

senthilsnc83@gmail.com (Senthil)

*Corresponding author

To cite this article:

Senthil, Nithya, Suryadevi, David Chandrakumar. (Alpha, Beta)-Normal and Skew n-Normal Composite Multiplication Operator on Hilbert Spaces. *International Journal of Discrete Mathematics*. Vol. 4, No. 1, 2019, pp. 45-51. doi: 10.11648/j.dmath.20190401.17

Received: February 15, 2019; Accepted: March 19, 2019; Published: May 6, 2019

Abstract: In this paper, the condition under which composite multiplication operators on Hilbert spaces become skew n-normal operators, (Alpha, Beta)-normal, parahyponormal and quasi-parahyponormal have been obtained in terms of radon-nikodym derivative.

Keywords: Composite Multiplication Operator, Conditional Expectation, Aluthge Transformation, Skew n-Normal Operator, Parahyponormal

1. Introduction

Let (X, \sum, μ) be a σ -finite measure space. Then a mapping T from X into X is said to be a measurable transformation if $T^{-1}(E) \in \sum$ for every $E \in \sum$. A measurable transformation T is said to be non-singular if $\mu(T^{-1}(E)) = 0$ whenever $\mu(E) = 0$. If T is non-singular then the measure μT^{-1} defined as $\mu T^{-1}(E) = \mu(T^{-1}(E))$ for every $E \in \sum$, is an absolutely continuous measure on \sum with respect to μ . Since μ is a σ -finite measure, then by the Radon-Nikodym theorem, there exists a non-negative function f_0 in $L^1(\mu)$ such that $\mu T^{-1}(E) = \int_E f_0 d\mu$ for every $E \in \sum$. The function f_0 is called the Radon-Nikodym derivative of μT^{-1} with respect to μ .

Every non-singular measurable transformation T from X into itself induces a linear transformation C_T on $L^p(\mu)$ defined as $C_T f = f \circ T$ for every f in $L^p(\mu)$. In case C_T is continuous from $L^p(\mu)$ into itself, then it is called a

composition operator on $L^p(\mu)$ induced by T . We restrict our study of the composition operators on $L^2(\mu)$ which has Hilbert space structure. If u is an essentially bounded complex-valued measurable function on X , then the mapping M_u on $L^2(\mu)$ defined by $M_u f = u \cdot f$, is a continuous operator with range in $L^2(\mu)$. The operator M_u is known as the multiplication operator induced by u . A composite multiplication operator is linear transformation acting on a set of complex valued \sum measurable functions f of the form

$$M_{u,T}(f) = C_T M_u(f) = (u \circ T)(f \circ T)$$

where u is a complex valued, \sum measurable function. In case $u=1$ almost everywhere, $M_{u,T}$ becomes a composition operator, denoted by C_T .

In the study considered is the using conditional expectation of composite multiplication operator on L^2 -spaces. For each $f \in L^p(X, \sum, \mu)$, $1 \leq p \leq \infty$, there exists a unique $T^{-1}(\sum)$ -measurable function $E(f)$ such that

$$\int_A g f d\mu = \int_A g E(f) d\mu$$

for every $T^{-1}(\sum)$ -measurable function g , for which the left integral exists. The function $E(f)$ is called the conditional expectation of f with respect to the subalgebra $T^{-1}(\sum)$. As an operator of $L^p(\mu)$, E is the projection onto the closure of range of T and E is the identity on $L^p(\mu)$, $p \geq 1$ if and only if $T^{-1}(\sum) = \sum$. Detailed discussion of E is found in [1-4].

1.1. (Alpha, Beta)--Normal Operator [13]

An operator T is called (α, β) -normal operator if $\alpha^2 T^* T \leq T T^* \leq \beta^2 T^* T$, ($0 \leq \alpha \leq 1 \leq \beta$)

1.2. Skew n-Normal Operator [12]

An operator T is called skew n -normal operator if $(T^n T^*) T = T (T^* T^n)$, for all natural number n .

1.3. p-Hyponormal Operator [15]

An operator T is called p -hyponormal operator if $(T^* T)^p \geq (T T^*)^p$, for $(0 < p < \infty)$

2. Related Work in the Field

The study of weighted composition operators on L^2 spaces was initiated by R. K. Singh and D. C. Kumar [5]. During the last thirty years, several authors have studied the properties of various classes of weighted composition operator. Boundedness of the composition operators in $L^p(\sum)$, ($1 \leq p < \infty$) spaces, where the measure spaces are σ -finite, appeared already in [6]. Also boundedness of weighted operators on $C(X, E)$ has been studied in [7]. Recently S. Senthil, P. Thangaraju and D. C. Kumar have proved several theorems on n -normal, n -quasi-normal, k -paranormal, and (n, k) paranormal of composite multiplication operators on L^2 spaces [8-11]. In this paper we investigate composite

$$(i) M_{u,T}^* M_{u,T} f = u^2 f_0 f$$

$$(ii) M_{u,T} M_{u,T}^* f = (u^2 \circ T) (f_0 \circ T) E(f)$$

$$(iii) M_{u,T}^n(f) = (C_T M_u)^n(f) = u_n (f \circ T^n), u_n = (u \circ T) (u \circ T^2) (u \circ T^3) \dots (u \circ T^n)$$

$$(iv) M_{u,T}^* f = (u f_0)(E(f) \circ T^{-1})$$

$$(v) M_{u,T}^{*n} f = u f_0 (E(u f_0) \circ T^{-(n-1)}) (E(f) \circ T^{-n})$$

$$\text{where } E(u f_0) \circ T^{-(n-1)} = (E(u f_0) \circ T^{-1}) (E(u f_0) \circ T^{-2}) \dots (E(u f_0) \circ T^{-(n-1)})$$

$$(vi) |M_{u,T}| = u \sqrt{f_0} f$$

$$(vii) |M_{u,T}^*| = (\sqrt{u^2 f_0 \circ T}) (E(\sqrt{u^2 f_0 \circ T} f))$$

with the notation from Herron's proposition,

multiplication operators of (α, β) -normal operator and skew n -normal operator $L^2(\mu)$ -spaces.

3. Hyponormality for Composite Multiplication Operator

The results in the following proposition were proved in [12], as part of his doctoral dissertation.

3.1. Proposition [3]

Let $E = E(\cdot \setminus A)$ and let φ be a non-negative F measurable function.

Define the positive operator P_φ by $P_\varphi f = \varphi E(\varphi f)$.

$$\text{Let } \hat{\varphi} = \frac{\varphi}{(E(\varphi^2))^{\frac{1}{4}}}. \text{ Then } P_{\hat{\varphi}}^{\frac{1}{2}} = P_{\hat{\varphi}}.$$

Define the operator R_φ by $R_\varphi f = E(\varphi f)$. Then

$$\|R_\varphi\| = \left\| \sqrt{E(\varphi^2)} \right\|_\infty.$$

In [3], has proved the following lemma, as noted for any non-negative function f ,

$$\sup \text{port } f \subset \sup \text{port } E(f^r) \text{ for any } r > 0$$

3.2. Lemma [14]

Let α and β be non-negative functions, with $S = \sup \text{port } \alpha$. Then the following are equivalent:

$$\text{For every } f \in L^2(\mu) \int_X \alpha |f|^2 d\mu \geq \int_X |E(\beta f \setminus A)|^2 d\mu$$

$$\sup \text{port } \beta \subset S \text{ and } E\left(\frac{\beta^2}{\alpha} \chi_S \setminus A\right) \leq 1 \text{ almost everywhere.}$$

3.3. Proposition

Let the composite multiplication operator $M_{u,T} \in B(L^2(\mu))$. Then for $u \geq 0$

$$|M_{u,T}^*| = P_{\sqrt{u^2 f_0} \circ T} = P_v, \text{ where } v = \frac{\sqrt{(u^2 f_0) \circ T}}{(E((u^2 f_0) \circ T))^{\frac{1}{4}}}$$

Theorem 3.1

Let the composite multiplication operator $M_{u,T} \in B(L^2(\mu))$ with weight $u \geq 0$ and S be the support of f_0 :

$$|M_{u,T}| \geq |C| \text{ if and only if } u^2 \geq 1$$

$$|C| \geq |M_{u,T}^*| \text{ if and only if } \sup \text{port } v = \sup \text{port } \sqrt{f_0}$$

$$\text{where } v = \frac{\sqrt{(u^2 f_0) \circ T}}{(E((u^2 f_0) \circ T))^{\frac{1}{4}}}, E \left(\frac{(u^2 f_0) \circ T}{(E((u^2 f_0) \circ T))^{\frac{1}{2}}} \frac{1}{\sqrt{f_0}} \chi_S \right) \leq 1 \text{ almost everywhere.}$$

Proof:

Since $|M_{u,T}|$ and $|C|$ are multiplication operators, we need only compare their symbols. After squaring, we get

$$|M_{u,T}| \geq |C| \text{ if and only if } u \sqrt{f_0} \geq \sqrt{f_0}$$

$$|M_{u,T}| \geq |C| \text{ if and only if } u^2 f_0 \geq f_0 \text{ Because } f_0 > 0 \text{ almost everywhere,}$$

we obtain (i).

$$\text{As for this f, } |C| \geq |M_{u,T}^*|$$

$$\forall f, \int \sqrt{f_0} |f|^2 d\mu \geq \langle |M_{u,T}^*| f, f \rangle$$

$$|M_{u,T}^*| = P_v, \text{ where } v = \frac{\sqrt{(u^2 f_0) \circ T}}{(E((u^2 f_0) \circ T))^{\frac{1}{4}}} = \int v E(v f) \overline{f} d\mu$$

However,

$$\int v E(v f) \overline{f} d\mu = \langle E(v f), v f \rangle = \|E(v f)\|^2 = \int |E(v f)|^2 d\mu$$

Since $\sup \text{port } v = \sup \text{port } \sqrt{f_0}$, the desired conclusion follows from lemma 3.2.

Theorem 3.5

Let the composite multiplication operator $M_{u,T} \in B(L^2(\mu))$. Then $M_{u,T}$ is p-hyponormal if and only if $u f_0 > 0, u f_0 \circ T > 0$

$$\text{and } E \left(\frac{1}{u^{2p} f_0} \right) \leq \frac{1}{u^{2p} f_0 \circ T}.$$

Proof:

$$\text{Here } \left\langle (M_{u,T}^* M_{u,T})^p f, f \right\rangle = \int_X u^{2p} f_0^p |f|^2 d\mu$$

$$\text{and } \left\langle (M_{u,T} M_{u,T}^*)^p f, f \right\rangle = \int_X \left| E \left((u^p f_0^{\frac{p}{2}}) \circ T f \right) \right|^2 d\mu$$

$$\text{This implies } \int_X u^{2p} f_0^p |f|^2 d\mu \geq \int_X \left| E \left((u^p f_0^{\frac{p}{2}}) \circ T f \right) \right|^2 d\mu$$

Since by lemma 2.2, for every $f \in L^2(\mu)$

$$\Rightarrow \sigma((u^p f_0^{\frac{p}{2}}) \circ T) \subset \sigma(u^p f_0^p) \text{ and } E \left[\frac{(u^{2p} f_0^p) \circ T}{u^{2p} f_0^p} \right] \leq 1$$

$$\Rightarrow E \left[\frac{1}{u^{2p} f_0^p} \right] \leq \frac{1}{(u^{2p} f_0^p) \circ T} \text{ if } (u^{2p} f_0^p) \circ T > 0 \text{ and } u^{2p} f_0^p > 0.$$

4. Parahyponormal for Composite Multiplication Operator

Mahmound M. Kutkut [16], has proved that an operator A is parahyponormal if and only if $(AA^*)^2 - 2CA^*A + C^2 \geq 0$ for all real C . In an analogous manner, we derive some characterization of parahyponormal and quasi-parahyponormal composite multiplication operator on L^2 -spaces.

Theorem 4.1

Let the composite multiplication operator $M_{u,T} \in B(L^2(\mu))$. Then $M_{u,T}$ is parahyponormal if and only if

$$u^4 \circ T \cdot f_0^2 \circ T E(f) - 2Cu^2 f_0 f + C^2 \geq 0 \text{ almost everywhere, for all } C \geq 0$$

Proof:

Suppose $M_{u,T}$ is parahyponormal. Then $(M_{u,T} M_{u,T}^*)^2 - 2C M_{u,T}^* M_{u,T} + C^2 \geq 0$ for all $C \geq 0$.

This implies that

$$\langle ((M_{u,T} M_{u,T}^*)^2 - 2C M_{u,T}^* M_{u,T} + C^2) f, f \rangle \geq 0 \text{ for all } f \in L^2(\mu)$$

By the proposition 3.3 we get,

$$\int_E \{ (u^2 \circ T \cdot f_0 \circ T E(f))^2 - 2Cu^2 f_0 f + C^2 \} d\mu \geq 0 \text{ for every } E \in \Sigma.$$

$$\Leftrightarrow u^4 \circ T \cdot f_0^2 \circ T E(f) - 2Cu^2 f_0 f + C^2 \geq 0 \text{ almost everywhere, for all } C \geq 0$$

Theorem 4.2

Let the composite multiplication operator $M_{u,T} \in B(L^2(\mu))$. Then $M_{u,T}$ is M-parahyponormal if and only if

$$m^2 u^4 \circ T \cdot f_0^2 \circ T E(f) - 2Cu^2 f_0 f + C^2 \geq 0 \text{ almost everywhere, for all } C \geq 0.$$

Proof:

Suppose $M_{u,T}$ is M-parahyponormal.

Then $m^2 (M_{u,T} M_{u,T}^*)^2 - 2C M_{u,T}^* M_{u,T} + C^2 \geq 0$ for all $C \geq 0$.

This implies that

$$\langle (m^2 (M_{u,T} M_{u,T}^*)^2 - 2C M_{u,T}^* M_{u,T} + C^2) f, f \rangle \geq 0 \text{ for all } f \in L^2(\mu)$$

By the proposition 3.3 we get,

$$\int_E \{ m^2 (u^2 \circ T \cdot f_0 \circ T E(f))^2 - 2Cu^2 f_0 f + C^2 \} d\mu \geq 0 \text{ for every } E \in \Sigma.$$

$$\Leftrightarrow m^2 u^4 \circ T \cdot f_0^2 \circ T E(f) - 2Cu^2 f_0 f + C^2 \geq 0 \text{ almost everywhere, for all } C \geq 0$$

Theorem 4.3

Let the composite multiplication operator $M_{u,T} \in B(L^2(\mu))$. Then $M_{u,T}$ is M^* -parahyponormal if and only if

$$m^2 u^2 f_0 E(u^2 f_0) \circ T^{-1} f - 2Cu^2 \circ T \cdot f_0 \circ T E(f) + C^2 \geq 0 \text{ almost everywhere, for all } C \geq 0.$$

Proof:

Suppose $M_{u,T}$ is M^* -parahyponormal.

Then $m^2 M_{u,T}^{*2} M_{u,T}^2 - 2C M_{u,T} M_{u,T}^* + C^2 \geq 0$ for all $C \geq 0$.

This implies that

$$\langle (m^2 M_{u,T}^{*2} M_{u,T}^2 - 2C M_{u,T} M_{u,T}^* + C^2) f, f \rangle \geq 0 \text{ for all } f \in L^2(\mu)$$

By the proposition 3.3 and $M_{u,T}^{*2} M_{u,T}^2 f = u^2 f_0 E(u^2 f_0) \circ T^{-1} f$ $u \geq 0$ we get,

$$\int_E \{ m^2 u^2 f_0 E(u^2 f_0) \circ T^{-1} f - 2Cu^2 \circ T \cdot f_0 \circ T E(f) + C^2 \} d\mu \geq 0 \text{ for every } E \in \Sigma.$$

$$\Leftrightarrow m^2 u^2 f_0 E(u^2 f_0) \circ T^{-1} f - 2C u^2 \circ T \cdot f_0 \circ T E(f) + C^2 \geq 0 \text{ almost everywhere, for all } C \geq 0$$

5. Quasi-Parahyponormal for Composite Multiplication Operator

By result of [16] an operators A on H is quasi-parahyponormal if and only if $(A^2 A^{*2})^2 - 2C(AA^*)^2 + C^2 \geq 0$ for all C . In an analogous manner, we derive the characterization of quasi-parahyponormal composite multiplication operator on L^2 -spaces.

Theorem 5.1

Let the composite multiplication operator $M_{u,T} \in B(L^2(\mu))$. Then $M_{u,T}$ is quasi-parahyponormal if and only if $u^2 \circ T \cdot u^4 \circ T^2 \cdot f_0^2 \circ T^2 \cdot (E(u f_0))^2 \circ T \cdot E(f) - 2C u^4 \circ T \cdot f_0^2 \circ T \cdot E(f) + C^2 \geq 0$ almost everywhere, for all $C \geq 0$

Proof:

Suppose $M_{u,T}$ is quasi-parahyponormal.

Then $(M_{u,T}^2 M_{u,T}^{*2})^2 - 2C(M_{u,T} M_{u,T}^*)^2 + C^2 \geq 0$ for all $C \geq 0$.

This implies that

$$\left\langle ((M_{u,T}^2 M_{u,T}^{*2})^2 - 2C(M_{u,T} M_{u,T}^*)^2 + C^2) f, f \right\rangle \geq 0 \text{ for all } f \in L^2(\mu)$$

By the proposition 3.3 and $M_{u,T}^2 M_{u,T}^{*2} f = u \circ T \cdot u^2 \circ T^2 \cdot f_0 \circ T^2 \cdot E(u f_0) \circ T \cdot E(f)$

$$\int_E \left\{ (u \circ T \cdot u^2 \circ T^2 \cdot f_0 \circ T^2 \cdot E(u f_0) \circ T \cdot E(f))^2 - 2C(u^2 \circ T \cdot f_0 \circ T \cdot E(f))^2 + C^2 \right\} d\mu \geq 0 \text{ for every } E \in \Sigma$$

$$\Leftrightarrow u^2 \circ T \cdot u^4 \circ T^2 \cdot f_0^2 \circ T^2 \cdot (E(u f_0))^2 \circ T \cdot E(f) - 2C u^4 \circ T \cdot f_0^2 \circ T \cdot E(f) + C^2 \geq 0$$

almost everywhere, for all $C \geq 0$

Theorem 5.2

Let the composite multiplication operator $M_{u,T} \in B(L^2(\mu))$. Then $M_{u,T}$ is M-quasi-parahyponormal if and only if $m^2 u^2 \circ T \cdot u^4 \circ T^2 \cdot f_0^2 \circ T^2 \cdot (E(u f_0))^2 \circ T \cdot E(f) - 2C u^4 \circ T \cdot f_0^2 \circ T \cdot E(f) + C^2 \geq 0$ almost everywhere, for all $C \geq 0$

Proof:

Suppose $M_{u,T}$ is M-quasi-parahyponormal.

Then $m^2 (M_{u,T}^2 M_{u,T}^{*2})^2 - 2C(M_{u,T} M_{u,T}^*)^2 + C^2 \geq 0$ for all $C \geq 0$.

This implies that

$$\left\langle (m^2 (M_{u,T}^2 M_{u,T}^{*2})^2 - 2C(M_{u,T} M_{u,T}^*)^2 + C^2) f, f \right\rangle \geq 0 \text{ for all } f \in L^2(\mu)$$

By the proposition 3.3 and $M_{u,T}^2 M_{u,T}^{*2} f = u \circ T \cdot u^2 \circ T^2 \cdot f_0 \circ T^2 \cdot E(u f_0) \circ T \cdot E(f)$

$$\int_E \left\{ m^2 (u \circ T \cdot u^2 \circ T^2 \cdot f_0 \circ T^2 \cdot E(u f_0) \circ T \cdot E(f))^2 - 2C(u^2 \circ T \cdot f_0 \circ T \cdot E(f))^2 + C^2 \right\} d\mu \geq 0 \text{ for every } E \in \Sigma$$

$$\Leftrightarrow m^2 u^2 \circ T \cdot u^4 \circ T^2 \cdot f_0^2 \circ T^2 \cdot (E(u f_0))^2 \circ T \cdot E(f) - 2C u^4 \circ T \cdot f_0^2 \circ T \cdot E(f) + C^2 \geq 0 \text{ almost everywhere, for all } C \geq 0$$

6. Skew n-normal and (Alpha, Beta)-normal Composite Multiplication Operator

Theorem 6.1

Let the composite multiplication operator $M_{u,T} \in B(L^2(\mu))$. Then $M_{u,T}$ is skew n-normal if and only if

$$u_n (u^2 \circ T^n) (f_0 \circ T^n) = (u^2 \circ T) (f_0 \circ T) (E(u_n) \circ T) \text{ almost everywhere.}$$

Proof:

$M_{u,T}$ is skew n-normal. Then

$$\begin{aligned} (M_{u,T}^n M_{u,T}^{*n}) M_{u,T} f &= (M_{u,T}^n M_{u,T}^{*n}) (u f) \circ T \\ &= M_{u,T}^n (u f_0 E((u f) \circ T) \circ T^{-1}) \end{aligned}$$

$$\begin{aligned}
&= M_{u,T}^n u^2 f_0 f \\
&= u_n(u^2 f_0 f) \circ T^n \\
&= u_n(u^2 \circ T^n)(f_0 \circ T^n)(f \circ T^n)
\end{aligned}$$

Also,

$$\begin{aligned}
M_{u,T}(M_{u,T}^* M_{u,T}^n) f &= M_{u,T} M_{u,T}^* (u_n(f \circ T^n)) \\
&= M_{u,T} u f_0 E(u_n(f \circ T^n)) \circ T^{-1} \\
&= M_{u,T} u f_0 E(u_n)(f \circ T^{n-1}) \\
&= u(u f_0 E(u_n)(f \circ T^{n-1})) \circ T \\
&= (u^2 \circ T)(f_0 \circ T) E(u_n) \circ T (f \circ T^n)
\end{aligned}$$

$$\Leftrightarrow u_n(u^2 \circ T^n)(f_0 \circ T^n) = (u^2 \circ T)(f_0 \circ T) (E(u_n) \circ T) \text{ almost everywhere.}$$

Theorem 6.2

Let the composite multiplication operator $M_{u,T} \in B(L^2(\mu))$. Then $M_{u,T}^*$ is skew n-normal if and only if $u^2 f_0^2 = (u^2 \circ T^{-(n-1)})(f_0 \circ T^{-(n-1)})$ almost everywhere.

Proof:

$M_{u,T}^*$ is skew n-normal. Then

$$\begin{aligned}
M_{u,T}^*(M_{u,T} M_{u,T}^{*n}) f &= M_{u,T}^* M_{u,T} \left(u f_0 \left(E(u f_0) \circ T^{-(n-1)} \right) \left(E(f) \circ T^{-n} \right) \right) \\
&= M_{u,T}^* \left(u \left(u f_0 \left(E(u f_0) \circ T^{-(n-1)} \right) \left(E(f) \circ T^{-n} \right) \right) \right) \circ T \\
&= M_{u,T}^* \left((u^2 \circ T)(f_0 \circ T) (E(u f_0) \circ T^{-(n+2)})(E(f) \circ T^{-(n-1)}) \right) \\
&= u f_0 E \left[\left((u^2 \circ T)(f_0 \circ T) (E(u f_0) \circ T^{-(n+2)})(E(f) \circ T^{-(n-1)}) \right) \right] \circ T^{-1} \\
&= u^3 f_0^2 (E(u f_0) \circ T^{-(n-1)})(E(f) \circ T^{-n})
\end{aligned}$$

Also,

$$\begin{aligned}
(M_{u,T}^{*n} M_{u,T}) M_{u,T}^* f &= M_{u,T}^{*n} M_{u,T} \left(u f_0 E(f) \circ T^{-1} \right) \\
&= M_{u,T}^{*n} \left((u^2 \circ T)(f_0 \circ T) E(f) \right) \\
&= u f_0 E(u f_0) \circ T^{-(n-1)} (u^2 \circ T^{-(n-1)})(f_0 \circ T^{-(n-1)}) E(f) \circ T^{-n}
\end{aligned}$$

$$\Leftrightarrow u^2 f_0^2 = (u^2 \circ T^{-(n-1)})(f_0 \circ T^{-(n-1)}) \text{ almost everywhere.}$$

Theorem 6.2

Let the composite multiplication operator $M_{u,T} \in B(L^2(\mu))$. Then $M_{u,T}$ is (α, β) -normal if and only if $\alpha^2 u^2 f_0 f \leq (u^2 f_0) \circ T E(f) \leq \beta^2 u^2 f_0 f$ almost everywhere.

Proof:

$M_{u,T}$ is (α, β) -normal. Then it is easy to check,

$$M_{u,T}^* M_{u,T} f = u^2 f_0 f, \quad M_{u,T} M_{u,T}^* f = (u^2 \circ T)(f_0 \circ T) E(f)$$

By definition,

$$\alpha^2 u^2 f_0 f \leq (u^2 f_0) \circ T E(f) \leq \beta^2 u^2 f_0 f \text{ almost everywhere.}$$

8. Conclusion

In the study of p-hyponormal operator, the Aluthge transform is a very useful tool. We investigate some basic properties of such operators and study the relation among skew n-normal operators, (α, β) -normal operator, parahyponormal and quasi-parahyponormal composite multiplication operators on $L^2(\mu)$ -space. In future try to generalize the composite multiplication operator on Poisson weighted sequence spaces.

Acknowledgements

We would like to thank the reviewers for carefully reading manuscript and for their constructive comments. I want to thank Professor Dr. R. David Chandrakumar, Department of Mathematics, Vickram College of Engineering for his support and encouragement during preparation of the paper.

References

- [1] Campbell J & Jamison J, On some classes of weighted composition operators, Glasgow Math. J. vol. 32, pp. 82-94, (1990).
- [2] Embry Wardrop M & Lambert A, Measurable transformations and centred composition operators, Proc. Royal Irish Acad, vol. 2(1), pp. 23-25 (2009).
- [3] Herron J, Weighted conditional expectation operators on Hilbert spaces, UNC charlotte doctoral dissertation.
- [4] Thomas Hoover, Alan Lambert and Joseph Quinn, The Markov process determined by a weighted composition operator, Studia Mathematica, vol. XXII (1982).
- [5] Singh RK & Kumar DC, Weighted composition operators, Ph.D. thesis, Univ. of Jammu (1985).
- [6] Singh RK, Composition operators induced by rational functions, Proc. Amer. Math. Soc., vol. 59, pp. 329-333(1976).
- [7] Takagi H & Yokouchi K, Multiplication and Composition operators between two Hilbert spaces, Contem. Math., vol. 232, pp. 321-338 (1999).
- [8] Panaiyappan S & Senthil kumar D, Parahyponormal and M-parahyponormal composition operators, Acta Ciencia Indica, vol. XXVIII (4) (2002).
- [9] Senthil S, Thangaraju P & Kumar DC, n-normal and n-quasi-normal composite multiplication operator on Hilbert spaces, Journal of Scientific Research & Reports, vol. 8(4), pp. 1-9 (2015).
- [10] Senthil S, Thangaraju P & Kumar DC, k-*paranormal, k-quasi-*paranormal and (n,k)-quasi-*paranormal composite multiplication operator on Hilbert spaces, British Journal of mathematics and computer science, BJMCS20166 (2015).
- [11] Senthil S, Thangaraju P & Kumar DC, Composite multiplication operator on Hilbert spaces of vector valued functions, International research Journal of Mathematical Sciences, vol. 4 (2), pp. 1(2015).
- [12] Shaakir LK & Abdulwahid ES, Skew n-normal operators, Aust. J. of basic and appl. sci, Vol-8(16), pp. 340-344 (2014).
- [13] Moslehian MS, On (Alpha, Beta)-normal operators in Hilbert spaces, IMAGE 39, Problem 39-4 (2007).
- [14] Burnap C, Jung I & Lambert A, Separating partial normality classes with composition operators, J. Operator Theory Vol. 53(2), pp. 381-397 (2005).
- [15] Fujii M & Nakatsu Y, On subclasses of hyponormal operators, Proc. Japan Acad. Ser. A. Math. Sci. vol. (51), pp. 243-246 (1975).
- [16] Kutkut & Mahmoud M, On the class of parahyponormal operators, J. Math Sci, (Calcutta), vol. 4(2), pp. 73-88 (1993).