

Adaptation Verification of Irrigated Cotton Cultivars for High Productivity and Economic Advantage for Large Scale Production at Tendaho Sugar Estate, Ethiopia

Zinaw Dilnesaw¹, Welday Gebreegziabher^{1, *}, Tesfay Hailu¹, Fikadu Abdisa¹, Abiy Getaneh¹, Mohamed Berhie²

¹Sugar Corporation, Research and Development Center, Wonji, Ethiopia

²Afar Region Pastoralists and Semi-pastoralist Agricultural Research, Semera, Ethiopia

Email address:

weldaygebre@gmail.com (Welday Gebreegziabher)

*Corresponding author

To cite this article:

Zinaw Dilnesaw, Welday Gebreegziabher, Tesfay Hailu, Fikadu Abdisa, Abiy Getaneh, Mohamed Berhie. Adaptation Verification of Irrigated Cotton Cultivars for High Productivity and Economic Advantage for Large Scale Production at Tendaho Sugar Estate, Ethiopia.

Ecology and Evolutionary Biology. Vol. 8, No. 3, 2023, pp. 67-73. doi: 10.11648/j.eeb.20230803.13

Received: June 17, 2023; **Accepted:** July 31, 2023; **Published:** September 8, 2023

Abstract: Adaptation performance field experiment on irrigated cotton cultivars for high productivity for large scale production and economic advantage were conducted in RCB Design by replicating four times to identify the high yielding with high quality cotton cultivar using surface irrigation at Tendaho sugar estate. All collected data were analyzed using GenStat 17th Edition software and mean comparisons among treatment means were made by LSD (5%). Cotton quality comparisons were also done based on the national quality measurement standards and the cotton quality laboratory analysis was done in Cotton Central accredited laboratory of Ethiopia under Ethiopian Textile Development and Research Institute. Significant difference ($p < 0.05$) among the tested cotton cultivars were reported for stand count (SC), number of unopened balls per plant and percent of 65 percent opened ball. However, there were no significant variability for total cotton yield (Y (kg/ha), lint yield and moisture adjusted yield (MoiAd_Y_Qt_Ha) at ($p < 0.05$) significance level. From the candidate cultivars, woyito and worer shown first and second high cotton productivity with 32 and 31.1 qt/ha cotton yield respectively followed by Stam-9-A as third with 30 qt/ha. In the other case, Stam-9-A and Delta ranked first for the major cotton quality standard measures. From this, it can be concluded that candidate genotype Stam-9-A which is third in productivity and first in cotton quality could have better advantage for the Estate. Therefore, based on the current adaptation trail result candidate cultivar Stam-9-A is recommended for commercial production in the study farm area.

Keywords: Adaptation, Cotton, Cultivar, Quality, Yield

1. Introduction

Agriculture is the 'engine for growth' in Africa with its immense land area covering 3 billion ha has 1.3 billion ha of agricultural land out of which only 252 million ha (19.36%) is arable [1]. With subsistence agriculture practiced by majority small holder farmers, yield gaps are high and poor soils, amongst other constraints add to the difficulties for sustainable farming and incomes. Average real per capital income in 2010 was \$688 (in constant 2000 US\$) compared to \$1717 in the rest of the developing world [2]. Sub-Saharan Africa has experienced encouraging economic growth

averaging about 4.5 per cent with some non-oil-exporting countries reaching an average of more than eight per cent [3]. Despite this impressive economic performance, agricultural transformation has been slow and growth rather sluggish.

Ethiopia as part of the region relies much on agriculture, although considerable and dynamic efforts made towards increasing agricultural production; the country has yet to go a long way to secure self-sufficiency in strategic food crops consequently, the country is obliged to import large quantities of wheat and other grains. Even though much of the irrigable lands are not yet used for various reasons and numerous reports indicated Ethiopia has excellent agro ecology as a

result have good agricultural potential that would allow to produce surplus agricultural products but yet the country dependent merely on rain-fed agriculture through harnessing its fertile and irrigable land in the lowland areas.

Cotton is more important than the various synthetic fibers, and it is grown all over the world in about 80 countries [4]. Currently, it is an important cash crop especially for a number of developing countries at local and national levels [5]. Ethiopia is one of the African countries that produce and export cotton [6]. The country is also believed to be one of the origins of cotton and its cultivation as it has a unique history in the country's agriculture because there are indigenous diploid ($2n = 26$) cultivated and wild species of cotton [7]. The indigenous cultivated species include *G. arboreum* and *G. herbaceum*. The distribution of the wild species of the B genome (*G. anomalum* subsp. *se-marensis*) and those of the E genome (*G. somalense*, *G. bricchettii* and *G. benadirensis*) have been documented. Ethiopia's major potential cotton growing areas include Omo-Ghibe, WabiShebele, Awash, Baro-Akobo, Blue Nile, and Tekeze river basins [8].

The basic conditions required for the effective production of cotton include a long frost-free period, a temperature range of 18–32°C and 600–1200 mm of water over the growing cycle, which typically lasts 125–175 days [9]. Therefore, Ethiopia has a very good cotton-growing condition and a large amount of land potentially suitable for cotton production [8]. The country possesses 2,697,640 hectares of land suitable for growing cotton; an area that equals the cotton land in Pakistan which is the world's fourth largest producer, but recently cultivated only 3% of this area for cotton production [10]. As a result, the amount of cotton produced in the country is small and the current domestic cotton production is much lower than the potential [11].

Ethiopia's dream is to become a middle income economy by 2025 by realizing an average annual economic growth of 10% through building a modern and productive agricultural sector, strengthening the industrial base and growing exports. Ethiopia's cotton production, which has remained fairly flat in recent years, was estimated at 40,000 metric tons 2014/15 and expands slightly in 2015/16 to 43,500 metric tons. Meantime, consumption is forecast to outstrip domestic cotton production in large part due to the demands of the rapidly-expanding textile/apparel industry. Thus, it is anticipated that imports, will fill this gap. Imports are estimated at 8,000 metric tons in 2014/15 and 12,000 metric

tons in 2015/16 [12].

Currently the Government of Ethiopia has started several development projects in every sector of the economy. As a part of this effort since 2010 onwards the Ethiopia Sugar Corporation is working to move the nation reach among the top ten sugar producing countries of the world by 2025 (GTP2), hence the corporation is developing irrigation infrastructures for about 420, 000 ha of land in Omo, Awash, Beless and Tekeze rivers basins (GTP-1) by establishing ten new sugar factories in different parts of the country. Among the newly established sugar estates, Tendaho has about 50,000ha farm land size. However, due to several reasons much of the irrigable land was not yet cultivated by sugarcane.

The global practices indicate that most of the sugar producing countries such as India, Thailand, Australia, South Africa and Brazil are running their sugar industries with complementary crops and livestock's enterprises. Therefore, Ethiopian Sugar Corporation is utilizing all efforts on harmonizing crop development with sugarcane and established a wing tasked with crop, horticulture and livestock production to enhance product diversification [13].

Even if EIAR has a role to coordinate national cotton research program in the country [14], most of the proposed large fertile irrigable low land areas are owned by sugar estates so that they have not been addressed by the national agricultural research systems in developing improved crop varieties yet. Therefore, it seems essential to undertake a specific variety adaptation trial at each location and generate scientific information on Ecological Suitability, specific Adaptability and economic Profitability of producing irrigated cotton at Tendaho Sugar Factory. Therefore, the objective of this verification trial was to evaluate Ecological Suitability, Adaptation and Profitability of irrigated cotton thereby, to identify high yielding cotton varieties better adapted to Tendaho sugar estate.

2. Material and Methods

2.1. Description of the Study Area

Tendaho Sugarcane Estate is found in Afar Regional State of Ethiopia located at 41°3'E longitude and 11°50' N latitude, 374 m above sea level elevation, receiving annual rainfall of about 200 mm with mean minimum and maximum temperature of 16.8°C and 38.8°C, respectively.

Table 1. Monthly Mean Maximum, Mean Minimum and Mean Average Temperature of Tendaho Sugarcane Estate (January, 1963 – December, 2018).

| Temperature (°C) | JAN | Feb | Mar | Apr | May | June | July | Aug | Sep | Oct | Nov | Dec | Mean |
|------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| Mean Maximum | 33.0 | 35.9 | 38.4 | 39.9 | 40.9 | 41.8 | 42.8 | 41.5 | 39.8 | 40.2 | 37.1 | 34.8 | 38.8 |
| Mean Minimum | 12.4 | 13.9 | 15.0 | 17.1 | 18.3 | 20.1 | 20.5 | 20.3 | 20.1 | 16.6 | 13.9 | 13.0 | 16.8 |
| Mean Average | 22.7 | 24.9 | 26.7 | 28.5 | 29.6 | 30.9 | 31.6 | 30.9 | 30.0 | 28.4 | 25.5 | 23.9 | 27.8 |

2.2. Methods and Materials Used in the Experiment

Five candidate Cotton genotypes; Delta, Inosa, Werer, Weyto and Stam -9-A developed in Worer Agricultural Center and recommended to agro ecologies of lower awash

valley and similar warm climates low altitude, were used for the trial at Tendaho Sugar Estate Dubti section. The seed source was Ethiopian Institute of Agricultural Research Worer Research center.

The candidate Cotton cultivars were evaluated in RCBD

with four replications using furrow irrigation at Tendaho Sugar Estate, Dubti site. The plot size of the trial was 10 m by 10 m (100 m²). The trial was carried out using furrow irrigation during June 2018 to end of October 2018 following recommended agronomic practices. Data were collected for major agronomic growth parameters, yield and yield components, cotton quality parameters and standard quality measurements parameters were collected based on ETDRI procedures.

2.3. Analysis of Variance

The data of different traits were collected and statistically analyzed using GenStat 17th Edition Software. Analysis of Variance for RCBD was computed. Mean comparisons among treatment means were also analyzed by LSD methods at 5% significance level.

The RCBD design analysis of variance was used to derive variance components as structured and stated by Cochran and Cox [15].

RCBD ANOVA was computed using the following model:

$$Y_{ij} = \mu + r_j + g_i + e_{ij}$$

Where, Y_{ij} = the response of trait Y in the i^{th} genotype and the j^{th} replication

μ = the grand mean of trait Y

r_j = the effect of the j^{th} replication

g_i = the effect of the i^{th} genotype

e_{ij} = experimental error effect

2.4. Profitability Analysis

2.4.1. Sources of Data

In this study primary and secondary sources of data were used. Cotton cultivars of farm production data were used as primary data and published and unpublished sources of data were used as secondary data source.

2.4.2. Data Analysis

In this study descriptive statistics were implemented as a tool for data analysis.

Table 2. Cotton Quality Standard.

| No | Descriptor | Grade A | Grade B | Grade C |
|----|---------------------------------|------------------|------------------|-----------------------|
| 1 | Moisture% | <8% | <8% | <8% |
| 2 | Micronier value* | 3.5 _ 4.2 | 4.3 _ 4.9 | 3.2 _ 3.4 and 5 _ 5.2 |
| 3 | Maturity | > 85% | 81 _ 84% | 75 _ 80% |
| 4 | UHML (mm)* | 28.5mm and above | 27mm _ 28.4mm | 25mm _ 26.9mm |
| 5 | UI%* | > 83% | 81 _ 82% | 76 _ 80% |
| 6 | SF% | <10% | 11 _ 20% | 21 _ 32 |
| 7 | Strength (g/tex)* | >29GPT | 26 _ 28.9GPT | 25 _ 25.9GPT |
| 8 | Color grade | 11 _ 1 to 21 _ 4 | 31 _ 1 to 31 _ 4 | 41 _ 1 to 51 _ 4 |
| 9 | Trash content (%) | < 3.5 | 3.5 _ 4.5% | 4.6 _ 5% |
| 10 | Average number of sticky points | 0 _ 10 | 11 _ 20 | 21 _ 32 |

Source: Ethiopian Textile Research and Development Institute Grading /ETRD IQG/

3. Result and Discussion

3.1. Variance

As indicated in Table 3, there were significant variation ($P < 0.05$) among the treatments for stand count (SC), number

of unopened balls per plant and percent of 65 percent opened ball. Stand count (SC) is one of the major characters of cotton that determine yield potential and has direct and significant relationship with number of effective balls per plant and total plot yield.

Table 3. NOVA, LSD and CV.

| Source of variation | d. f. | s. s. | m. s. | v. r. | F pr. | l. s. d. | cv% |
|---------------------|-------|--------|--------|-------|--------|----------|------|
| Stand ct | 4 | 231.2 | 57.8 | 5.3 | 0.011* | 5.086 | 7.8 |
| Residual | 12 | 130.8 | 10.9 | | | | |
| PH | 4 | 1665.9 | 416.5 | 2.81 | 0.074 | 18.74 | 14.7 |
| Residual | 12 | 1776.4 | 148 | | | | |
| Peaking_date_First | 4 | 25.7 | 6.425 | 2.69 | 0.083 | 2.383 | 1.3 |
| Residual | 12 | 28.7 | 2.392 | | | | |
| No_Un_Oped | 4 | 61.648 | 15.412 | 3.75 | 0.033* | 3.123 | 35 |
| Residual | 12 | 49.308 | 4.109 | | | | |
| No_Open_Ball | 4 | 12.24 | 3.06 | 0.28 | 0.885 | 5.085 | 24.6 |
| Residual | 12 | 130.73 | 10.89 | | | | |
| Initial_Squaring | 4 | 8.7 | 2.175 | 1.05 | 0.423 | 2.219 | 4.9 |
| Residual | 12 | 24.9 | 2.075 | | | | |
| Initial_Flowering | 4 | 5.3 | 1.325 | 1.67 | 0.22 | 1.371 | 2.2 |

| Source of variation | d. f. | s. s. | m. s. | v. r. | F pr. | l. s. d. | cv% |
|-----------------------|-------|---------|----------|-------|--------|----------|------|
| Residual | 12 | 9.5 | 0.7917 | | | | |
| Initial_Boll_opening | 4 | 77.3 | 19.325 | 2.56 | 0.093 | 4.236 | 3.1 |
| Residual | 12 | 90.7 | 7.558 | | | | |
| First_picking_kg | 4 | 275.61 | 68.9 | 2.3 | 0.118 | 8.43 | 22.2 |
| Residual | 12 | 359.38 | 29.95 | | | | |
| Emergence | 4 | 2.7 | 0.675 | 0.42 | 0.791 | 1.954 | 20.1 |
| Residual | 12 | 19.3 | 1.608 | | | | |
| Average_boll_wt_gram | 4 | 1.5778 | 0.3944 | 1.97 | 0.163 | 0.689 | 7.7 |
| Residual | 12 | 2.403 | 0.2002 | | | | |
| %65%_Boll_opening | 4 | 253.7 | 63.425 | 6.54 | 0.005* | 4.796 | 2.9 |
| Residual | 12 | 116.3 | 9.692 | | | | |
| %50%_Flowering | 4 | 2.3 | 0.575 | 0.26 | 0.898 | 2.289 | 2.8 |
| Residual | 12 | 26.5 | 2.208 | | | | |
| %30bolls_weight_kg | 4 | 0.00147 | 0.000368 | 3.13 | 0.056 | 0.0167 | 6.2 |
| Residual | 12 | 0.00141 | 0.000118 | | | | |
| %30_bolls_weight_gram | 4 | 1408.5 | 352.1 | 1.95 | 0.166 | 20.69 | 7.7 |
| Residual | 12 | 2165 | 180.4 | | | | |

3.2. Yield Comparison

As shown in the result, there was variability among the studied candidate genotypes for the rest twelve agronomic traits even though there were statistically not significant. There were no significant variability for total cotton yield (Y

(kg/ha), lint yield and moisture adjusted yield (MoiAd_Y_Qt_Ha) (table 4). There was no significant variability among tested cultivars for cotton yield as indicated in table (4 and 5) in variance analysis and mean comparison.

Table 4. ANOVA for Cotton Yield in kg/ha, qt/ha and moisture adjusted yield qt/ha.

| | Source of variation | d. f. | s. s. | m. s. | v. r. | F pr. |
|---------------|---------------------|---------|----------|---------|-------|-------|
| Y (kg/ha) | Rep stratum | 3 | 2382217 | 794072 | 1.78 | |
| | Rep.*Units* stratum | | | | | |
| | Tret | 4 | 4098963 | 1024741 | 2.3 | 0.118 |
| | Residual | 12 | 5344695 | 445391 | | |
| | Total | 19 | 11825876 | | | |
| FY (qt/Ha) | Source of variation | d. f. | s. s. | m. s. | v. r. | F pr. |
| | Rep stratum | 3 | 238.22 | 79.41 | 1.78 | |
| | Rep.*Units* stratum | | | | | |
| | Tret | 4 | 409.9 | 102.47 | 2.3 | 0.118 |
| | Residual | 12 | 534.47 | 44.54 | | |
| MoiAd_Y_Qt_Ha | Source of variation | d. f. | s. s. | m. s. | v. r. | F pr. |
| | Rep stratum | 3 | 201.63 | 67.21 | 1.78 | |
| | Rep.*Units* stratum | | | | | |
| | Tret | 4 | 346.94 | 86.73 | 2.3 | 0.118 |
| | Residual | 12 | 452.38 | 37.7 | | |
| Total | 19 | 1000.94 | | | | |

Table 5. Cotton Yield mean comparison.

| Treatments | Y (kg/ha) | Tret | FY (qt/Ha) | Tret | MoiAd_Y_Qt_Ha |
|------------|-----------|------|------------|-------|---------------|
| weyto | 3474 | a | Weyto | 34.74 | a |
| werer | 3383 | a | Werer | 33.83 | a |
| Stam -9-A | 3261 | a | Stam -9-A | 32.61 | a |
| Inosa | 2507 | a | Inosa | 25.07 | a |
| delta | 2415 | a | Delta | 24.15 | a |

Even though there was no significant variability among studied cultivars for cotton yield, but there was high variation

in magnitude for cotton yield productivity (table 6). Hence, candidate cultivar Woyto recorded the first higher adjusted

cotton yield (32 Qt/ha), followed by Werer (31.1 Qt/ha) second and Stam-9-A (30 Qt/ha) as a third.

Table 6. Mean, Grand mean, l. s. d. and CV for cotton yield.

| | delta | Inosa | werer | weyto | Stam -9-A | Grand | l. s. d. | CV |
|---------------|-------|-------|-------|-------|-----------|-------|----------|------|
| Y (kg/ha) | 2415 | 2507 | 3383 | 3474 | 3261 | 3008 | 1028.2 | 22.2 |
| FY (qt/Ha) | 24.2 | 25.1 | 33.8 | 34.7 | 32.6 | 30.08 | 10.28 | 22.2 |
| MoiAd Y Qt Ha | 22.2 | 23.1 | 31.1 | 32 | 30 | 27.68 | 9.46 | 22.2 |

3.3. Cotton Quality

As indicated in the material and method section, cotton quality comparison were done based on the national quality measurement standard of the country and the cotton quality laboratory analysis was done in Cotton Central accredited laboratory of Ethiopia under Ethiopian Textile Development and Research Institute.

The quality analysis were done for Moisture%, Micronier value, Maturity, UHML (mm), UI%, SF%, Strength (g/tex), Color grade, Trash content (%) and Average number of sticky points. As indicated in table 7, candidate cultivars quality grade to Moisture% recorded small variation within the range of the standard for example werer, weyto and Stam -9-A scored grade A for 6.85%, 6.28% and 6.73% respective value of moisture measurement.

Similarly, candidate cultivars did not show significant

variation for Micronier value and all candidate cultivars scored grade C quality with indicated values in table 7. All of the candidate cultivars scored quality grade A for a parameter Maturity; while Delta and Stam-9-A scored quality grade A for UHML (mm) parameter whereas Inosa, werer and Woyito scored B as indicated in table 7. Except woyito (B) all candidate cultivars scored quality grade A for UI%* parameter.

All candidate cultivars scored quality A for SF% and Color grade parameters as shown in table 7. Four candidate cultivars scored quality grade B for Strength (g/tex) quality parameter but Stam-9-A scored A. while, delta, Inosa, werer and weyto scored B for Trash Content (%) but Stam -9-A scored lower quality grade C. Similarly as indicated in (table 7) candidate cultivar delta scored A for Av No sticky points, grade score B for Inosa and Stam-9-A while C for werer as indicated.

Table 7. Cotton quality ETRDI Laboratory Result Report.

| Treatments | Moisture% | Micronier value* | Maturity | UHML (mm) | UI%* |
|------------|-----------|------------------|-------------|-------------------|---------------------|
| delta | 6.83 | 5.64 | 0.88 | 28.58 | 83.8 |
| ETRDI QG | A | C | A | A | A |
| Inosa | 6.73 | 5.85 | 0.88 | 27.57 | 83.2 |
| ETRDI QG | A | C | A | B | A |
| werer | 6.85 | 5.32 | 0.87 | 28.26 | 83.9 |
| ETRDI QG | A | C | A | B | A |
| weyto | 6.28 | 5.24 | 0.87 | 27.09 | 82.6 |
| ETRDI QG | A | C | A | B | B |
| Stam -9-A | 6.73 | 5.22 | 0.88 | 30.07 | 84.6 |
| ETRDI QG | A | C | A | A | A |
| Treatments | SF% | Strength (g/tex) | Color grade | Trash Content (%) | Av No sticky points |
| delta | 8.23 | 26.08 | 11-1 | 0.375 | 7.5 |
| ETRDI QG | A | B | A | B | A |
| Inosa | 8.45 | 27.38 | 11-2 | 0.375 | 14 |
| ETRDI QG | A | B | A | B | B |
| werer | 8.33 | 26.23 | 11-1 | 0.426 | 21.25 |
| ETRDI QG | A | B | A | B | C |
| weyto | 8.9 | 26.65 | 11-1 | 0.358 | 14.75 |
| ETRDI QG | A | B | A | B | B |
| Stam -9-A | 7.3 | 29.6 | 11-2 | 0.492 | 16 |
| ETRDI QG | A | A | A | C | B |

Quality Summary

Table 8. Cotton quality test comparison summary.

| Treatments | Quality Rank |
|------------|-----------------|
| Stam -9-A | 1 st |
| delta | 1 st |
| Inosa | 3 nd |
| werer | 3 nd |
| weyto | 5 th |

As shown in the table 8 Delta and Stam-9-A ranked first for the major cotton quality standard measures. But woyito and werer shown first and second high cotton productivity with 32 and 31.1 qt cotton yield quintal per hectare respectively followed third by Stam-9-A 30 qt/ha.

4. Conclusion and Recommendation

Cultivars Woyito and Worer scored first and second high cotton yield as compared to the others with 32 and 31.1 qt/ha respectively followed by Stam -90A with 30 qt/ha ranked third. As indicated there with respect of cotton quality comparison, Stam-9-A and Delta scored first in quality parameters. The result and discussion indicated cultivars

variance analysis shown non-significant variability for Yield and related parameters and cotton grade quality is a major determinant factor for produce market. Therefore, we can conclude that candidate genotype Stam-9-A which is third in productivity and first in cotton quality could have better advantage for the Estate. Hence, based on the current adaptation trail result candidate Stam-9-A is recommended for commercial production in Tendaho Dubti farm area.

Appendix

Mean Comparison.

Table A1. Mean LSD Multiple range test.

| Treatments | Stand ct | comp | PH | comp | 1 st Peaking date | comp | No Un Open | comp | No Open Ball | comp |
|------------|----------|------|-------|------|------------------------------|------|------------|------|--------------|------|
| delta | 38.75 | b | 88.82 | a | 122.2 | a | 9.05 | a | 12.88 | a |
| Inosa | 44 | a | 93 | a | 121.8 | ab | 5.65 | b | 13.62 | a |
| werer | 37.75 | b | 67.17 | b | 120 | ab | 5.75 | b | 14.82 | a |
| weyto | 46.25 | a | 78.4 | ab | 119.2 | b | 4.15 | b | 12.65 | a |
| Stam -9-A | 44.75 | a | 86.32 | ab | 121.5 | ab | 4.35 | b | 13.03 | a |

| Treatments | Initial Squaring | comp | Initial Flowering | comp | Initial Bo opening | comp | First Picking /kg | comp | Emergence | comp |
|------------|------------------|------|-------------------|------|--------------------|------|-------------------|------|-----------|------|
| delta | 29.5 | a | 41.5 | ab | 88 | ab | 19.81 | a | 6.5 | a |
| Inosa | 29.5 | a | 42.25 | a | 91.75 | a | 20.56 | a | 5.75 | a |
| werer | 31 | a | 41 | ab | 88.25 | ab | 27.74 | a | 6 | a |
| weyto | 29.25 | a | 41.25 | ab | 85.75 | b | 28.49 | a | 6.5 | a |
| Stam -9-A | 29.25 | a | 40.75 | b | 89.5 | ab | 26.74 | a | 6.75 | a |

| Treatments | Average bo wt/gm | comp | %65%_Boll opening | comp | %50%_Flowering | comp | %30bolls Weight/kg | comp | %30_bolls Weight/gram | comp |
|------------|------------------|------|-------------------|------|----------------|------|--------------------|------|-----------------------|------|
| delta | 5.315 | ac | 111.5 | a | 53 | a | 0.16 | b | 159.5 | ac |
| Inosa | 5.912 | abc | 112 | a | 52.25 | a | 0.1775 | a | 177.3 | abc |
| werer | 6.062 | a | 107.2 | ab | 52 | a | 0.1825 | a | 181.8 | ab |
| weyto | 5.697 | abc | 102.5 | b | 52.25 | a | 0.17 | ab | 170.9 | abc |
| Stam -9-A | 6.06 | ab | 110.8 | a | 52.25 | a | 0.1825 | a | 181.8 | a |

References

- [1] FAO, 2011. The state of food and agriculture. Women in Agriculture. Closing the Gender gap for Development.
- [2] Feeding Africa, 2015. Action Plan for African Agricultural Transformation, United Nations Economic Commission of Africa, Conference Background Paper, Cereal Crops, Harold Macauley, Director General of Africa Rice and Tabo Ramadjita, ICRISAT, 21-23 October 2015, Dakar Senegal.
- [3] Nicole Mason et al, 2012. Wheat Consumption in Sub-Saharan Africa: Trends, Drivers, and Policy Implications, MSU International Development Working Paper 127, December 2012.
- [4] Acquaah, G. 2007. Principles of Plant Genetics and Breeding, 2nd edition. Garsington Road, Oxford OX42 DQ, UK.
- [5] Baffes, J. 2004. Cotton market setting, trade policies, and issue. World Bank Policy Research Working Paper 3218; Washington DC.
- [6] Melesse, Z., Mohammed, A., Mezgebu, A. and Habtamu, M. 2019. Cotton production and marketing trend in Ethiopia: Cotton production and marketing trend in Ethiopia, Cogent Food and Agriculture, 5: 1, 1691812.
- [7] Merdasa Balcha, Nurhussien Seid, Yonas Bekele, Michael Kebede, Donis Gurmessa, Zemedkun Alemu, Samuel Damtew, Arkebe G/egziabher, Seleshi Getahun, Fikremariam Tegegn, Mekashaw Arega and Tamiru Dejen. 2022. Cotton Production Guideline in Ethiopia. Werer Agricultural Research Center (WARC), Ethiopian Institute of Agricultural Research (EIAR), Addis Ababa, Ethiopia. ISBN978-99944-3-509-8 ©EIAR, 2022 Webpage: <http://www.eiar.gov.et> Addis Ababa, Ethiopia.
- [8] Alebel, B., Firew, B., Berihu, A. and Mezgebe, M. 2014. An institutional assessment of the cotton and sugarcane commodities in Ethiopia. The climate change perspective. Addis Ababa, Ethiopia: Ethiopian Development Research Institute (EDRI). [Google Scholar].
- [9] FAO, 2012. Crop yield response to water. FAO irrigation and drainage paper No. 66. Rome.
- [10] EIA (Ethiopian Investment Agency). 2012. Investment opportunity profile for cotton production and ginning In Ethiopia. Updated Report on Cotton Ginning. Addis Ababa, Ethiopia. doi: 10.1094/PDIS-11-11-0999-PDN [Crossref]. [Google Scholar].
- [11] Bosena, T., Bekabil, F., Gebremedhin, B. and Hoekstra D. 2011. Factors affecting cotton supply at the farm level in Metema district of Ethiopia. Journal of Agriculture, Biotechnology and Ecology, 4 (1): 41-51.

- [12] Global Agricultural Information Network, GAIN, 2012. "Ban on Cotton Exports Lifted," GAIN Report Number: ET 1204, Addis Ababa.
- [13] Adaptation and Promotion of Selected Cereal and Forage Crops in Newly Established Sugar Estates Ethiopian Agricultural Research Council Secretariat, July 2016, Addis Ababa, Ethiopia.
- [14] Bedane G. and Arkebe G. 2019. Cotton production potential areas, production trends, research status, gaps and future directions of cotton improvement in Ethiopia. *Greener Journal of Agricultural Sciences* 9 (2): 163-170, <http://doi.org/10.15580/GJAS.2019.2.040619064>.
- [15] Cochran, W. G. and Cox, G. M., 1957. *Experimental Designs*. New York: Wiley.