



Biological Agents of Bioremediation: A Concise Review

Karabi Biswas, Dipak Paul, Sankar Narayan Sinha*

Environmental Microbiology Research Laboratory, Department of Botany, University of Kalyani, Kalyani, West Bengal, India

Email address:

biswaskarabi@yahoo.in (K. Biswas), dipak23paul@yahoo.com (D. Paul), sinhasn62@yahoo.co.in (S. N. Sinha)

To cite this article:

Karabi Biswas, Dipak Paul, Sankar Narayan Sinha. Biological Agents of Bioremediation: A Concise Review. *Frontiers in Environmental Microbiology*. Vol. 1, No. 3, 2015, pp. 39-43. doi: 10.11648/j.fem.20150103.11

Abstract: Due to intensive agriculture, rapid industrialization and anthropogenic activities have caused environmental pollution, land degradation and increased pressure on the natural resources and contributing to their adulteration. Bioremediation is the use of biological organisms to destroy, or reduce the hazardous wastes on a contaminated site. Bioremediation is the most potent management tool to control the environmental pollution and recover contaminated soil. Use of biological materials, coupled to other advanced processes is one of the most promising and inexpensive approaches for removing environmental pollutants. Bioremediation technology is a beneficial alternative which leads to degrade of pollutants. This article presents the important biological organisms used in bioremediation technologies.

Keywords: Bioremediation, Biological Organisms, Environmental Pollution

1. Introduction

Heavy metals and organic pollutants are considered to be a significant environmental issue for human health. The contamination of soil and water bodies by organic pollutants and toxic metals has been increased for the past few years due to industrialization, intensive agriculture and anthropogenic activity. Indiscriminate and unrestricted disposal of industrial effluents and urban city sewage into the aquatic environment has become an issue of major global concern [1-3]. Excess loading of unsafe and hazardous waste as led to shortage of clean and hygienic water and disruption of soil this limiting crop production [4].

Bioremediation processes are more attractive than physical and chemical techniques such as ion exchange, electrochemical treatment, reverse osmosis, evaporation, precipitation and sorption for heavy metal removal techniques for lower cost and higher efficiency at low metal concentrations. There are a number of biological materials that can be utilized to remove the hazardous metal from waste water such as bacteria, fungi and algae [5, 6].

Most of the advances in bioremediation have been realized through the assistance of the scientific areas of biochemistry, microbiology, analytical chemistry, molecular biology, environmental and chemical engineering, among others. These different fields, each with its own individual approach, have actively contributed to the development of bioremediation progress in recent years [7].

Now-a-days, the experience accumulated over the last

decades, has improved our understanding in many aspects of this multidisciplinary technology. The combined evaluation of the technical and non technical issues is an important step towards the successful application of environmental biotechnology in remediation. The states of the art of bioremediation technology as well as examples of more or less successful case studies are published in many books during the last two decades [8-11].

In this review, different techniques used in bioremediation and the ability of some bacteria, fungi and algae to remediate the organic and inorganic pollutants have been critically discussed.

2. Mechanism of Bioremediation

Bioremediation is a biological treatment system to destroy, or reduce the concentration of hazardous waste from a contaminated site. Thus some definitions restrict to the use of microbes only while others seem to incorporate all the biological entities such as plants (phytoremediation). Whatever, it can be define, in fact in nature the process of biological remediation involves both plants and microbes and rather the plant-microbe interaction in root zone has a very important role. pH is the important factor influencing the adsorption. Crist *et al.* [12] reported that with decreasing pH, the number of binding sites reduced and that pH increased during the metal ion uptake.

3. Agents of Bioremediation

Natural organisms, either indigenous or extraneous, are the important agents used for bioremediation [13]. The organisms vary, depending on the chemical properties of the polluting substances, and are to be chosen cautiously as they only sustain within a stipulated limit of chemical contaminants [13, 14]. The first patent for a biological remediation substance was recorded in 1974, was a strain of *Pseudomonas putida* capable of degrading petroleum [13]. Bioremediation can take place naturally or through intervention processes [15].

4. Bioremediation by Bacteria

The microbes have often been reported for the degradation of pesticides and hydrocarbons. A large number of bacteria utilize the contaminant as the sole carbon and energy sources. They are listed in Table 1 and Table 2.

Metals play important role in the life processes of microbes. Some metals such as chromium (Cr), calcium (Ca), magnesium (Mg), manganese (Mn), copper (Cu), sodium (Na), nickel (Ni) and zinc (Zn) are essential as micronutrients for various metabolic functions and for redox functions.

Table 1. Bioaccumulation and biotransformation of organic molecules by bacteria.

Bacteria	Toxic chemicals	References
<i>Bacillus</i> sp	Cresol, phenols, aromatics, long chain alkanes, phenol,	[16]
<i>Pseudomonas</i> sp	Benzene, anthracene, hydrocarbons, PCBs	[16, 17]
<i>Flavobacterium</i> sp	Aromatics	[18]
<i>Azotobacter</i> sp	Aromatics	[18]
<i>Xanthomonas</i> sp	Hydrocarbons, polycyclic hydrocarbons	[18, 19]
<i>Nocardia</i> sp	Hydrocarbons	[20]
<i>Streptomyces</i> sp	Phenoxyacetate, halogenated hydrocarbon, diazinon	[18]
<i>Mycobacterium</i> sp	Aromatics, branched hydrocarbons benzene, cycloparaffins	[21]

Table 2. Heavy metals utilizing bacteria.

Bacteria	Heavy metals	References
<i>Bacillus</i> sp	Cu, Zn	[22]
<i>Pseudomonas aeruginosa</i>	U, Cu, Ni, Cr	[23-25]
<i>Aerococcus</i> sp	Pb, Cr, Cd	[26]
<i>Aeromonas</i> sp	Cr	[25]
<i>Rhodopseudomonas palustris</i>	Pb	[27]
<i>Citrobacter</i> sp	Cd, U, Pb	[22, 28]

5. Bioremediation by Fungus

Fungi represent the promising group of microbes for biodegradation (Table 3 and Table 4). The ability of fungi, both yeasts and moulds, to convert a broad variety of

hazardous chemical substances has developed interest to use them in bioremediation [9]. Fungi can mineralize xenobiotic compounds to CO₂ and H₂O through their non-specific ligninolytic and highly oxidative enzyme system, which is also responsible for the degradation and decolorization of a wide range of dyes [29, 30].

Table 3. Bioaccumulation and biotransformation of organic molecules by fungi.

Fungi	Toxic chemicals	References
<i>Coprinellus radians</i>	PAHs, methylnaphthalenes, and dibenzofurans	[31]
<i>Marasmiellus troyanus</i>	Benzo [a] pirene	[32]
<i>Gloeophyllum trabeum</i>	1, 1, 1-trichloro-2, 2-bis (4-chlorophenyl) ethane (DDT)	[33]
<i>Pleurotus ostreatus</i>	Bisphenol A	[34]
<i>Fomitopsis pinicola</i>	1, 1, 1-trichloro-2, 2-bis (4-chlorophenyl) ethane (DDT)	[33]
<i>Penicillium simplicissimum</i>	Polyethylene	[35]
<i>Phanerochaete chrysosporium</i>	Polyvinylamine sulfonate anthrapyridone	[36]

Table 4. Heavy metals utilizing fungi.

Fungi	Heavy metals	References
<i>Rhizopus arrhizus</i>	Ag, Hg	[22]
<i>Stereum hirsutum</i>	Cd, Pb	[37, 38]

6. Bioremediation by Algae

Biodegradation of pesticides is determined by two groups of factors, the first relates to microbial consortium and the optimum condition for their survival and activity while the

second relates to the chemical structure of the pesticides. Factors related to microorganisms including the presence and number of appropriate microorganisms, the contact between microorganisms and the substrate (pesticide), pH, temperature, salinity, nutrients, light quality and intensity, available water, oxygen tension and redox potential, surface binding, presence of alternative carbon substrates and alternative electron acceptors. Kobayashi and Rittman [39] showed that not only microalgae have the ability to bioaccumulate pesticides, but also capable to biotransform

some of these environmental contaminants.

Table 5. Bioaccumulation and biotransformation of organic molecules by algae.

Algae	Elements	References
<i>Chlamydomonas</i> sp	Naphthalene	[39]
<i>Dunaliella</i> sp	Naphthalene, DDT	[39]
<i>Euglena gracilis</i>	DDT, Phenol	[39]
<i>Selenastrum capricornutum</i>	Benzene, toluene, chlorobenzene, 1, 2- dichlorobenzene, nitrobenzene naphthalene, 2, 6-dinitrotoluene, phenanthrene, di-n-butylphthalate, pyrene	[39]
<i>Chlorella</i> sp	Toxaphene	[39]
<i>Cylindrotheca</i> sp	DDT	[39]

The capability of algae to absorb hazardous metals has been known for many years. Algae have the ability to remove toxic heavy metals from the environment, which results in higher concentrations than those of the surrounding water [40]. Algae, related eukaryotic photosynthetic organisms, have exclusively developed the production of different peptides capable to attach heavy metals. These organometallic complexes are further separated inside vacuoles facilitating proper regulation of the heavy metal ions concentration of cytoplasm, thus neutralizing or preventing their toxic effect [41].

Table 6. Heavy metals utilizing algae.

Algae	Heavy metals	References
<i>Zooglea</i> sp	Co, Ni, Cd	[22]
<i>Phormidium valderium</i>	Cd, Co, Cu, Ni	[37, 38]
<i>Chlorella vulgaris</i>	Au, Cu, Ni, U, Pb, Hg, Zn	[22, 42]
<i>Volvariella volvacea</i>	Cu, Hg, Pb	[43, 44]
<i>Oscillatoria</i> sp	Ni, Cu, Co, Pb, Zn	[45]
<i>Tetraselmis chuii</i>	Cu	[46]
<i>Spirogyra hyalina</i>	Cd, Hg, Pb, As	[47]
<i>Chlorella pyrenoidosa</i>	Zn, Cu, As, Pb, Cd, Cr, Ni, Hg	[48]
<i>Lyngbya spiralis</i>	Cd, Pb, Hg	[49]

Table 7. Bioremediation by higher plant.

Plants	Contaminants	References
<i>Helianthus annuus</i>	Cu, Pb, EDTA	[50]
<i>Brassica juncea</i>	Pb, Cr, Ni, Zn	[51, 52]
<i>Perennial ryegrass</i>	Cd, Ag	[53]
<i>Alfa alfa</i>	Chlorinated aliphatics	[50]
<i>Nicotiana tabacum</i>	Se, Hg, Ti	[50]
<i>Tagetes patula</i>	Benzo [a] pyrene	[54]
<i>Arundo donax</i>	As	[55]
<i>Typha domingensis</i>	Al, Fe, Zn, Pb	[56]
<i>Solanum nigrum</i>	Cd	[57]
<i>Lemna minor</i>	Cd, Hg, Zn, Mn, Pb, Ag	[58]

7. Bioremediation by Higher Plants

Phytoremediation is well suited for applications in low-permeability soils, where most currently utilized technologies have a low level of success, as well as in combination with more conventional removal technologies (foam migration, electromigration, etc.). In appropriate situations, phytoremediation can be a substitute to the much harsher physical and chemical remediation technologies of thermal vaporization, solvent washing, incineration or other soil washing techniques, which essentially degrade the biological constituents of the soil and can intensely change

its physical and chemical properties as well, generating a comparatively nonviable solid waste. Phytoremediation actually improves the soil, leaving a better, effective, soil ecosystem at costs estimated at around one-tenth of those presently acquired methods.

8. Conclusion

Soil and water are being polluted by various organic and inorganic pollutants due to rapid industrialization and use of agrochemicals in imbalanced proportions. Restrictive and clean up measures to avert hazards from contaminated soil belong to the curative soil protection. Bioremediation is a unique and cost-effective technique for cleaning up pollution by intensifying the natural biodegradation processes. So developing an understanding of microbial and plant communities with their response to the natural environment and contaminants, elaborating the knowledge of the genetics of the microorganisms helps to increase capabilities to degrade pollutants and recovery of land and ground water.

References

- [1] Strong PJ, Burgess JE (2008) Treatment methods for wine related ad distillery wastewaters: a review. *Bioremediation Journal* 12 (2): 70-87.
- [2] Paul D, Sinha SN (2013) Assessment of various heavy metals in surface water of polluted sites in the lower stretch of river Ganga, West Bengal: a study for ecological impact. *Discovery Nature* 6 (14): 8-13.
- [3] Sinha SN, Paul D (2014) Biodegradation potential of some bacterial strains isolated from sewage water. *International Journal of Environmental Biology* 4 (2): 107-111.
- [4] Kamaludeen SPB, Arukumar KR, Avudainayagam S, Ramasamy K (2003) Bioremediation of chromium contaminated environments. *Indian Journal of Experimental Biology* 41 (9): 972-985.
- [5] Vieira RH, Volesky B (2000) Biosorption: a solution to pollution? *International Microbiology* 3 (1): 17-24.
- [6] Sinha SN, Paul D (2012) Detoxification of heavy metals by biosurfactants. *Bulletin of Environmental and Scientific Research* 1 (3-4): 1-3.
- [7] Sheehan D (1997) *Bioremediation Protocols*, Humana Press, Totowa, New Jersey.

- [8] Cookson JT (1995) Bioremediation Engineering. Design and Application, McGraw-Hill, New York.
- [9] Alexander M (1999) Biodegradation and Bioremediation, Academic Press, San Diego, CA.
- [10] Filler DM, Snape I, Barnes DL (2008) Bioremediation of Petroleum Hydrocarbons in Cold Regions, Cambridge University Press.
- [11] Neilson AH, Allard AS (2008) Environmental degradation and transformation of organic chemicals, CRC Press.
- [12] Crist RH, Martin JR, Carr D (1994) Interaction of metals and protons with algae. *Environmental Science and Technology* 28 (11): 1859- 1866.
- [13] Prescott LM, Harley JP, Klein DA (2002) Microbiology, 5th edition, McGrawHill, New York.
- [14] Dubey RC (2004) a text book of Biotechnology 3rd edition, S Chand and Company Ltd. New Delhi, India.
- [15] Agarwal SK (1998) Environmental Biotechnology (1st edition), APH Publishing Corporation, New Delhi, India.
- [16] Cybulski Z, Dzurla E, Kaczorek E, Olszanowski A (2003). The influence of emulsifiers on hydrocarbon biodegradation by Pseudomonadacea and Bacillaceae strains. *Spill Science and Technology Bulletin* 8 (5): 503–507.
- [17] Kapley A, Purohit HJ, Chhatre S, Shanker R, Chakrabarti T (1999) Osmotolerance and hydrocarbon degradation by a genetically engineered microbial consortium. *Bioresource Technology* 67 (3): 241-245.
- [18] Jogdand SN (1995). Environmental biotechnology, 1st edition, Himalaya Publishing House, Bombay, India.
- [19] Ijah UJJ (1998) Studies on relative capabilities of bacterial and yeast isolates from tropical soil in degrading crude oil. *Waste Management* 18 (5): 293–299.
- [20] Park AJ, Cha DK, Holsen TM (1998) Enhancing solubilization of sparingly soluble organic compounds by biosurfactants produced by *Nocardia erythropolis*. *Water Environment Research* 70 (3): 351– 355.
- [21] Lee S, Cutright TJ (1995) Bioremediation of polycyclic aromatic hydrocarbon-contaminated soil. U. S. Patent No. 5, 427, 944. Washington, DC: U. S. Patent and Trademark Office.
- [22] Rajendran P, Muthukrishnan J, Gunasekaran P (2003) Microbes in heavy metal remediation. *Indian journal of experimental biology* 41 (9): 935-944.
- [23] Sar P, D'Souza SF (2001) Biosorptive uranium uptake by *Pseudomonas* strain: characterization and equilibrium studies. *Journal of Chemical Technology and Biotechnology* 76 (12): 1286-1294.
- [24] Sar P, Kazy SK, Asthana RK, Singh SP (1999) Metal adsorption and desorption by lyophilized *Pseudomonas aeruginosa*. *International Biodeterioration and Biodegradation* 44 (2): 101-110.
- [25] Sinha SN, Biswas M, Paul D, Rahaman S (2011) Biodegradation potential of bacterial isolates from tannery effluent with special reference to hexavalent chromium. *Biotechnology Bioinformatics and Bioengineering* 1 (3): 381-386.
- [26] Sinha SN, Paul D (2014) Heavy metal tolerance and accumulation by bacterial strains isolated from waste water. *Journal of Chemical, Biological and Physical Sciences* 4 (1): 812- 817.
- [27] Sinha SN, Biswas K (2014) Bioremediation of lead from river water through lead-resistant purple-nonsulfur bacteria. *Global Journal of Microbiology and Biotechnology* 2 (1): 11-18.
- [28] Yan G, Viraraghavan T (2001) Heavy metal removal in a biosorption column by immobilized *M. rouxii* biomass. *Bioresource Technology* 78 (3): 243-249.
- [29] Fu Y, Viraraghavan T (2001) Fungal decolorization of dye wastewaters: a review. *Bioresource Technology* 79 (3): 251-262.
- [30] Wesenberg D, Kyriakides I, Agathos SN (2003) White-rot fungi and their enzymes for the treatment of industrial dye effluents. *Biotechnology Advances* 22 (1): 161-187.
- [31] Aranda E, Ullrich R, Hofrichter M (2010) Conversion of polycyclic aromatic hydrocarbons, methyl naphthalenes and dibenzofuran by two fungal peroxxygenases. *Biodegradation* 21 (2): 267–281.
- [32] Wunch KG, Alworth WL, Bennett JW (1999) Mineralization of benzo [a] pyrene by *Marasmiellus troyanus*, a mushroom isolated from a toxic waste site. *Microbiological Research* 154 (1): 75–79.
- [33] Purnomo AF, Kamei I, Kondo R (2008) Degradation of 1, 1, 1-trichloro-2, 2-bis (4-chlorophenyl) ethane (DDT) by brown-rot fungi. *Journal of Bioscience and Bioengineering* 105 (6): 614–621.
- [34] Hirano T, Honda Y, Watanabe T, Kuwahara M (2000) Degradation of bisphenol A by the lignin-degrading enzyme, manganese peroxidase, produced by the white-rot basidiomycete, *Pleurotus ostreatus*. *Bioscience, Biotechnology, and Biochemistry* 64 (9): 1958-1962.
- [35] Yamada-Onodera K, Mukumoto H, Katsuyama Y, Tani Y (2002) Degradation of long-chain alkanes by a polyethylene-degrading fungus, *Penicillium simplicissimum* YK. *Enzyme and Microbial Technology* 30 (6): 828-831.
- [36] Darah I, Ibrahim CO (1997) Lignin-degrading enzyme production by entrapped cells of *Phanerochaete chrysosporium* in submerged culture system. *Asia-Pacific Journal of Molecular Biology and Biotechnology* 5 (3): 143-154.
- [37] Gabriel J, Mokrejs M, Bily J, Rychlovsky P (1994) Accumulation of heavy metal by some Wood- rooting fungi. *Folia Microbiologica* 39 (2), 115-118.
- [38] Gabriel J, Kofronova O, Rychlovsky P, Krenzelok M (1996) Accumulation and effect of cadmium in the wood rotting basidiomycete, *Daedalea quercina*. *Bulletin of Environmental Contamination and Toxicology* 57 (3): 383-390.
- [39] Kobayashi H, Rittman BE (1982) Microbial removal of hazardous organic compounds. *Environment Science and Technology* 16 (3): 170A-183A.
- [40] Megharaja M, Ragusa SR, Naidu R (2003) Metal–algae interactions: implication of bioavailability. In: Naidu, R., Gupta, V. V. S. R., Rogers, S., Kookana, R. S., Bolan, N. S. and Adriano, D. C. (eds) *Bioavailability, Toxicity and Risk Relationships in Ecosystems*, Science Publishers, Enfield, New Hampshire pp. 109–144.

- [41] Cobbett C, Goldsbrough P (2002) Phytochelatin and metallothioneins: Roles in heavy metal detoxification and homeostasis. *Annual Review of Plant Biology* 53 (1):159–182.
- [42] Pearson RG (1969) Hard and soft acids and bases. *Survey of Progress in Chemistry* 5 (1): 1-52.
- [43] Purkayastha RP, Mitra AK (1992) Metal uptake by mycelia during submerged growth and by sporocarp of an edible fungus, *Volvariella volvacea*. *Indian Journal of Experimental Biology* 30 (12): 1184-1187.
- [44] Jagadevan S, Mukherji S (2004) Successful in-situ oil bioremediation programmes – key parameters. *Indian Journal of Biotechnology* 3 (4): 495–501.
- [45] Ajayan KV, Selvaraju M, Thirugnanamoorthy K (2011) Growth and heavy metals accumulation potential of microalgae grown in sewage wastewater and petrochemical effluents. *Pakistan Journal of Biological Sciences* 14 (16): 805-811.
- [46] Yılmaz AB, Işık O, Sayın S (2005) Bioaccumulation and toxicity of different copper concentrations in *Tetraselmis chuii*. *E. U. Journal of Fisheries and Aquatic Sciences* 22 (3-4): 297-304.
- [47] Kumar JIN, Oommen C (2012) Removal of heavy metals by biosorption using freshwater alga *Spirogyra hyalina*. *Journal of Environmental Biology* 33: 27-31.
- [48] Yao J, Li W, Xia F, Zheng Y, Fang C, Shen D (2012) Heavy metals and PCDD/Fs in solid waste incinerator fly ash in Zhejiang province, China: chemical and bio-analytical characterization. *Environmental Monitoring and assessment* 184 (6): 3711-3720.
- [49] Inthorn D, Sidtitoon N, Silapanuntakul S, Incharoensakdi A (2002) Sorption of mercury, cadmium and lead by microalgae. *Science Asia* 28: 253-261.
- [50] Terry N, Carlson C, Raab TK, Zayed AM (1992) Rates of selenium volatilization among crop species. *Journal of Environmental Quality* 21 (3): 341-344.
- [51] Pilon-Smith EAH, De Souza MP, Hong G, Amini A, Bravo RC, Payabyab ST, Terry N (1999) Selenium volatilization and accumulation by twenty aquatic plant species. *Journal of Environmental Quality* 28 (3): 1011-1017.
- [52] Heaton ACP, Rugh CL, Wang N, Meagher RB (1998) Phytoremediation of mercury- and methylmercury-polluted soils using genetically engineered plants. *Journal of Soil Contamination* 7 (4): 497-510.
- [53] Rugh CL, Gragson GM, Meagher RB, Merkle SA (1998) Toxic mercury reduction and remediation using transgenic plants with a modified bacterial gene. *Hortscience* 33 (4): 618-621.
- [54] Sun Y, Zhou Q, Xu Y, Wang L, Liang X (2011) Phytoremediation for co-contaminated soils of benzo [a] pyrene (B [a] P) and heavy metals using ornamental plant *Tagetes patula*. *Journal of Hazardous Materials* 186 (2): 2075-2082.
- [55] Mirza N, Mahmood Q, Pervez A, Ahmad R, Farooq R, Shah MM, Azim MR (2010) Phytoremediation potential of *Arundo donax* in arsenic-contaminated synthetic wastewater. *Bioresource Technology* 101 (15): 5815-5819.
- [56] Hegazy AK, Abdel-Ghani NT, El-Chaghaby GA (2011) Phytoremediation of industrial wastewater potentiality by *Typha domingensis*. *International Journal of Environmental Science and Technology* 8 (3): 639-648.
- [57] Ji P, Sun T, Song Y, Ackland ML, Liu Y (2011) Strategies for enhancing the phytoremediation of cadmium-contaminated agricultural soils by *Solanum nigrum* L. *Environmental Pollution* 159 (3): 762-768.
- [58] Ugya AY (2015) The Efficiency of *Lemna minor* L. in the phytoremediation of Romi stream: A case study of Kaduna Refinery and Petrochemical Company polluted stream. *Journal of Applied Biology and Biotechnology* 3 (1): 11-14.